arXiv:1106.4717v3 [hep-ph] 30 Sep 2011

KANAZAWA-11-11, KIAS-P11037

Two Component Dark Matters in $S_4 \times Z_2$ Flavor Symmetric Extra U(1) Model

Yasuhiro Daikoku[a1], Hiroshi Okada[b,c2] and Takashi Toma[a,d3]

[a] *Institute for Theoretical Physics, Kanazawa University, Kanazawa 920-1192, Japan*

[b] *Centre for Theoretical Physics, The British University in Egypt,*
El Sherouk City, Postal No, 11837, P.O. Box 43, Egypt

[c] *School of Physics, KIAS, Seoul 130-722, Korea*

[d] *Max-Planck-Institut für Kernphysik, Postfach 103980, 69029 Heidelberg, Germany*

Abstract

We study cosmic-ray anomaly observed by PAMELA based on E_6 inspired extra U(1) model with $S_4 \times Z_2$ flavor symmetry. In our model, the lightest flavon has very long lifetime of $\mathcal{O}(10^{18})$ second which is longer than the age of the universe, but not long enough to explain the PAMELA result $\sim \mathcal{O}(10^{26})$ sec. Such a situation could be avoidable by considering that the flavon is not the dominant component of dark matters and the dominant one is the lightest neutralino. With appropriate parameter set, density parameter of dark matter and over-abundance of positron flux in cosmic-ray are realized at the same time. There is interesting correlation between spectrum of positron flux and V_{MNS}. No excess of anti-proton in cosmic-ray suggests that sfermions are heavier than 4 TeV and the masses of the light Higgs bosons are degenerated.

---

[1] yasu_daikoku@yahoo.co.jp

[2] hokada@kias.re.kr

[3] t-toma@hep.s.kanazawa-u.ac.jp

1 Introduction

Standard Model (SM) is successful theory of gauge interactions, however Higgs sector is not examined well. Therefore mass matrices of leptons and quarks are not well understood. Many unsolved puzzles of SM are left in these sectors; that is, *e.g.*, why is the structure of mixing matrix of leptons (Maki-Nakagawa-Sakata matrix, V_{MNS}) very different from that of the mixing matrix of quarks (Cabibbo-Kobayashi-Maskawa matrix, V_{CKM}), especially why is the mixing angle θ_{23} maximal? Why is neutrino mass far smaller than those of other fermions? Why do generations exist?

We also find a problem in cosmology. In modern cosmology, the existence of the dark matter is clear. Recent cosmic-ray observation of PAMELA suggests that the dark matter decays mainly into leptons with very long lifetime [1][2][3]. Such a particle is not included in SM.

Separately from these puzzles, there is hierarchy problem why electroweak scale is much smaller than Planck scale. One of the solutions is to introduce supersymmetry (SUSY) [5]. However minimal supersymmetric standard model (MSSM) does not satisfy the solution, because we must fine-tune μ-parameter in superpotential of MSSM, which is much smaller than Planck scale in order to realize appropriate electroweak symmetry breaking. This is called μ-problem.

Another problem of MSSM is proton stability. The R-parity forbids baryon number violating trilinear terms in superptential, however does not forbid quartic terms like $E^cU^cU^cD^c, LQQQ$. Such interactions reduce the lifetime of proton to unacceptable level [6]. Therefore the R-parity does not help the explanation of proton stability. The problem of proton lifetime of supersymmetric model is one of the most essential point in understanding generation structure.

With the motivation to solve flavor puzzles and hierarchy problem, we introduce new three symmetries. At first, we introduce non-Abelian discrete flavor symmetry $S_4 \times Z_2$, in order to explain that the mixing angle θ_{23} is maximal [7][8][9][10][11]. Because V_{CKM} and V_{MNS} are very different, it is expected that the representations of quarks and leptons are also different. Next, we introduce $U(1)_X$ gauge symmetry which forbids μ-term [12]. Then, several new superfields must be introduced due to gauge anomaly cancellation condition; those are extra Higgs (H^U, H^D), singlet Higgs S and exotic quarks (g, g^c). The extra Higgs bosons couple only to leptons, which induce the difference between V_{CKM} and V_{MNS}. Moreover, the existence of exotic quarks is important to understand the meaning of generations. Finally we introduce $U(1)_Z$ gauge symmetry. Due to the anomaly cancellation condition, right-handed neutrino (RHN) superfield N^c is introduced, then the smallness of neutrino mass is realized by seesaw mechanism. The two new $U(1)$ gauge symmetries and standard model gauge symmetry $G_{SM} = SU(3)_c \times SU(2)_W \times U(1)_Y$ can be embedded in E_6 as $G_{SM} \times U(1)_X \times U(1)_Z \subset E_6$, then MSSM and new superfields consist **27** of E_6 representation. With appropriate assignment of superfields under the flavor symmetry, the stability of proton is realized, which plays the most important role in the flavor symmetry. Thus we can understand that the generation structure is the new system to stabilize proton [8].

The new symmetries which are introduced above may also solve dark matter problems. As three $U(1)$ gauge symmetries include R-parity, lightest supersymmetric particle (LSP) is a candidate for dark matter. The positron flux observed by PAMELA is produced by the field which induces RHN mass and decays into leptons [13]. In this paper, we show our model is consistent with experimental results of dark matter. At first, we define our model in section 2. The estimations of relic abundance of dark matter and positron flux are given in section 3. Finally, we give conclusion of our analysis in section 4.

2 $S_4 \times Z_2$ flavor symmetric extra U(1) model

2.1 Gauge symmetry

We extend the gauge symmetry from G_{SM} to $G_2 = G_{SM} \times U(1)_X \times U(1)_Z \subset E_6$, and add new superfields S, g, g^c, N^c which are embedded in **27** representation of E_6 with quark, lepton superfields Q, U^c, D^c, L, E^c and Higgs superfields H^U, H^D. In order to break $U(1)_Z$ gauge symmetry, we introduce G_{SM} singlet Φ and Φ^c. The gauge representations of these superfields are given in Table 1. After the gauge symmetry breaking, as the R-parity symmetry

$$R = \exp\left[\frac{i\pi}{20}(3x - 8y + 15z)\right] \tag{1}$$

	Q	U^c	E^c	D^c	L	N^c	H^D	g^c	H^U	g	S	Φ	Φ^c
$SU(3)_c$	3	3^*	1	3^*	1	1	1	3^*	1	3	1	1	1
$SU(2)_W$	2	1	1	1	2	1	2	1	2	1	1	1	1
$y = 6Y$	1	-4	6	2	-3	0	-3	2	3	-2	0	0	0
x	1	1	1	2	2	0	-3	-3	-2	-2	5	0	0
z	-1	-1	-1	2	2	-4	-1	-1	2	2	-1	8	-8
R	$-$	$-$	$-$	$-$	$-$	$-$	$+$	$+$	$+$	$+$	$+$	$+$	$+$
q_ψ	1	1	1	1	1	1	-2	-2	-2	-2	4	-2	2
q_χ	-1	-1	-1	3	3	-5	-2	-2	2	2	0	10	-10

Table 1: G_2 assignment of superfields. Where the x, y and z are charges of $U(1)_X$, $U(1)_Y$ and $U(1)_Z$, and Y is hypercharge. The extra U(1) charges x and z are given by $x = \frac{5}{4}q_\psi + \frac{1}{4}q_\chi$ and $z = -\frac{1}{4}q_\psi + \frac{3}{4}q_\chi$, where $q_\psi = 6\sqrt{\frac{2}{5}}Q_\psi$ and $q_\chi = 2\sqrt{6}Q_\chi$ are charges of two U(1)s in $SU(5) \times U(1)_\psi \times U(1)_\chi \subset SO(10) \times U(1)_\psi \subset E_6$.

remains unbroken, LSP is the candidate for dark matter. The invariant superpotential under the gauge symmetry G_2 is given by

$$\begin{aligned} W &= Y^U H^U Q U^c + Y^D H^D Q D^c + Y^E H^D L E^c \\ &+ \lambda S H^U H^D + k S g g^c \\ &+ Y^N H^U L N^c + Y^M \Phi N^c N^c \\ &+ M_\Phi \Phi^c \Phi + y_1 Q Q g + y_2 g^c U^c D^c + y_3 g E^c U^c + y_4 g^c L Q + y_5 g D^c N^c, \end{aligned} \tag{2}$$

where first line consists of trilinear terms in MSSM. Second line generates effective μ term $\lambda \langle S \rangle H^U H^D$ by radiative symmetry breaking of $U(1)_X$. Third line generates RHN mass term $Y^N \langle \Phi \rangle N^c N^c$ by radiative symmetry breaking of $U(1)_Z$ and gives small neutrino mass by seesaw mechanism. Fourth line consists of unwanted terms which cause the problems such that the mass term $M_\Phi \Phi \Phi^c$ prevents Φ, Φ^c from developing vacuum expectation values (VEVs) and the trilinear terms of exotic quarks destabilize proton. Note that Higgs superfields are extended to three generations. Generally, extra Higgs doublets cause the problem of flavor changing neutral currents (FCNCs).

2.2 Flavor symmetry

	Q_1	Q_2	Q_3	U_1^c	U_2^c	U_3^c	D_1^c	D_2^c	D_3^c
S_4	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**	**1**
Z_2	$+$	$+$	$+$	$+$	$+$	$+$	$+$	$+$	$+$

	E_1^c	E_2^c	E_3^c	L_i	L_3	N_i^c	N_3^c	H_i^D	H_3^D
S_4	**1**	**1**	**1′**	**2**	**1**	**2**	**1**	**2**	**1**
Z_2	$+$	$-$	$+$	$-$	$-$	$+$	$-$	$-$	$+$

	H_i^U	H_3^U	S_i	S_3	g_a	g_a^c	Φ_i	Φ_3	Φ_a^c
S_4	**2**	**1**	**2**	**1**	**3**	**3**	**2**	**1**	**3**
Z_2	$-$	$+$	$-$	$+$	$+$	$+$	$-$	$+$	$+$

Table 2: $S_4 \times Z_2$ assignment of superfields (Where the index i of the S_4 doublets runs $i = 1, 2$, and the index a of the S_4 triplets runs $a = 1, 2, 3$.)

In ordre to explain maximal mixing angle θ_{23}, we introduce $S_4 \times Z_2$ flavor symmetry. This symmetry solves the problems of superpotential defined in Eq.(2) at the same time. If we assign g, g^c, Φ^c to S_4 -triplets and the other superfields to singlets or doublets, then $M_\Phi, y_{1,\cdots,5}$ are eliminated. As a result, superpotential is given by

$$\begin{aligned} W' &= Y^U H^U Q U^c + Y^D H^D Q D^c + Y^E H^D L E^c \\ &+ \lambda S H^U H^D + k S g g^c \end{aligned}$$

$$\begin{aligned} &+ \quad Y^N H^U L N^c + Y^M \Phi N^c N^c \\ &+ \quad \frac{a}{M_P}\Phi\Phi\Phi^c\Phi^c \\ &+ \quad \frac{y'_1}{M_P^2}\Phi\Phi^c QQg + \frac{y'_2}{M_P^2}\Phi\Phi^c g^c U^c D^c + \frac{y'_3}{M_P^2}\Phi\Phi^c g E^c U^c \\ &+ \quad \frac{y'_4}{M_P^2}\Phi\Phi^c g^c LQ + \frac{y'_5}{M_P^2}\Phi\Phi^c g D^c N^c + \cdots, \end{aligned} \tag{3}$$

where the dots $\cdots$ are higher order terms. As the potential of Φ and Φ^c is lifted by non-renormalizable term $\Phi\Phi\Phi^c\Phi^c$, Φ and Φ^c have very large VEVs along the D-flat direction of $\langle\Phi\rangle = \langle\Phi^c\rangle$, where

$$V = \langle\Phi\rangle = \langle\Phi^c\rangle \sim \left(\frac{M_P m_{SUSY}}{a}\right)^{\frac{1}{2}} \sim 10^{11}\mathrm{GeV}\frac{1}{\sqrt{a}}\left(\frac{m_{SUSY}}{10\mathrm{TeV}}\right)^{\frac{1}{2}}. \tag{4}$$

From the constraints on the lifetimes of proton and exotic quarks (see appendix C), the condition

$$\frac{V^2}{M_P^2} \sim 10^{-12} \tag{5}$$

must be satisfied [14]. From this condition, $V \sim 10^{12}$GeV is required. This value is realized by potential minimum condition as Eq.(4), when we take $a \sim 10^{-2}$. The prediction for RHN mass is given by

$$M_R \sim 10^{12}\mathrm{GeV}, \tag{6}$$

which gives an appropriate neutrino mass. As the VEVs of Φ and Φ^c break not only $U(1)_Z$ but also $S_4 \times Z_2$, we call them flavons.

2.3 Maki-Nakagawa-Sakata matrix V_{MNS}

The maximal mixing angle θ_{23} of V_{MNS} is realized by the assignments that H^U, H^D, L, N^c are $\mathbf{2}+\mathbf{1}$ of S_4 and E^c is $\mathbf{1}+\mathbf{1}+\mathbf{1'}$ [7]. In order to reduce the number of parameters, we assign S and Φ to $\mathbf{2}+\mathbf{1}$. If we assign Q, U^c and D^c to S_4-singlets, then quarks do not couple to S_4 doublet Higgs and FCNC is suppressed. The flavor representations are given in Table 2. The leading order superpotential is given by

$$W_{S_4\times Z_2} = W_L + W_Q + W_H + W_g + W_\Phi, \tag{7}$$

$$\begin{aligned} W_L &= Y_2^N\left[H_1^U(L_1N_2^c + L_2N_1^c) + H_2^U(L_1N_1^c - L_2N_2^c)\right] \\ &+ \quad Y_3^N H_3^U L_3 N_3^c + Y_4^N L_3(H_1^U N_1^c + H_2^U N_2^c) \\ &+ \quad Y_1^E E_1^c(H_1^D L_1 + H_2^D L_2) + Y_2^E E_2^c H_3^D L_3 + Y_3^E E_3^c(H_1^D L_2 - H_2^D L_1) \\ &+ \quad \frac{1}{2}Y_1^M \Phi_3(N_1^cN_1^c + N_2^cN_2^c) + \frac{1}{2}Y_3^M\Phi_3 N_3^cN_3^c, \end{aligned} \tag{8}$$

$$W_Q = Y_{ij}^U H_3^U Q_i U_j^c + Y_{ij}^D H_3^D Q_i D_j^c \quad (i,j = 1,2,3), \tag{9}$$

$$\begin{aligned} W_H &= \lambda_1 S_3(H_1^U H_1^D + H_2^U H_2^D) + \lambda_3 S_3 H_3^U H_3^D \\ &+ \quad \lambda_4 H_3^U(S_1 H_1^D + S_2 H_2^D) + \lambda_5(S_1 H_1^U + S_2 H_2^U)H_3^D, \end{aligned} \tag{10}$$

$$W_g = k S_3(g_1 g_1^c + g_2 g_2^c + g_3 g_3^c), \tag{11}$$

$$\begin{aligned} W_\Phi &= \frac{a_1}{2M_P}\Phi_3^2[(\Phi_1^c)^2 + (\Phi_2^c)^2 + (\Phi_3^c)^2] \\ &+ \quad \frac{a_2}{2M_P}(\Phi_1^2 + \Phi_2^2)[(\Phi_1^c)^2 + (\Phi_2^c)^2 + (\Phi_3^c)^2] \\ &+ \quad \frac{a_3}{2M_P}\left\{2\sqrt{3}\Phi_1\Phi_2[(\Phi_2^c)^2 - (\Phi_3^c)^2] + (\Phi_1^2 - \Phi_2^2)[(\Phi_2^c)^2 + (\Phi_3^c)^2 - 2(\Phi_1^c)^2]\right\}. \end{aligned} \tag{12}$$

We give the parameter set to realize the maximal mixing angle of MNS matrix. We define non-negative VEVs as

$$\begin{aligned} &\langle H_1^U\rangle = \langle H_2^U\rangle = \frac{1}{\sqrt{2}}v_u, \quad \langle H_3^U\rangle = v'_u, \quad \langle H_1^D\rangle = \langle H_2^D\rangle = \frac{1}{\sqrt{2}}v_d, \quad \langle H_3^D\rangle = v'_d, \\ &\langle S_1\rangle = \langle S_2\rangle = \frac{1}{\sqrt{2}}v_s, \quad \langle S_3\rangle = v'_s, \end{aligned}$$

$$\langle\Phi_1\rangle = v_4, \quad \langle\Phi_2\rangle = \sqrt{v_1^2+v_2^2+v_3^2-v_4^2}, \quad \langle\Phi_3\rangle = V,$$
$$\langle\Phi_1^c\rangle = \sqrt{\frac{V^2}{3}+v_1^2}, \quad \langle\Phi_2^c\rangle = \sqrt{\frac{V^2}{3}+v_2^2}, \quad \langle\Phi_3^c\rangle = \sqrt{\frac{V^2}{3}+v_3^2} \quad (v_{1,2,3,4} \ll V). \tag{13}$$

In W_L, without loss of generality, we can define $Y_{2,4}^N, Y_{1,2,3}^E, Y_{1,3}^M$ to be real and define the phase of Y_3^N as $Y_3^N = |Y_3^N|e^{i\delta}$. We define mass parameters as

$$\begin{array}{lll} M_1 = Y_1^M V, & M_3 = Y_3^M V, & \\ m_2^\nu = Y_2^N v_u, & m_3^\nu = |Y_3^N| v_u', & m_4^\nu = Y_4^N v_u, \\ m_1^l = Y_1^E v_d, & m_2^l = Y_2^E v_d', & m_3^l = Y_3^E v_d. \end{array} \tag{14}$$

Using these parameters, the mass matrices of charged leptons and neutrinos are given by

$$M_l = \frac{1}{\sqrt{2}}\begin{pmatrix} m_1^l & 0 & -m_3^l \\ m_1^l & 0 & m_3^l \\ 0 & \sqrt{2}m_2^l & 0 \end{pmatrix}, \quad M_D = \frac{1}{\sqrt{2}}\begin{pmatrix} m_2^\nu & m_2^\nu & 0 \\ m_2^\nu & -m_2^\nu & 0 \\ m_4^\nu & m_4^\nu & \sqrt{2}e^{i\delta}m_3^\nu \end{pmatrix},$$
$$M_R = \begin{pmatrix} M_1 & 0 & 0 \\ 0 & M_1 & 0 \\ 0 & 0 & M_3 \end{pmatrix}. \tag{15}$$

The following neutrino mass matrix is generated through the seesaw mechanism

$$M_\nu = M_D M_R^{-1} M_D^t = \begin{pmatrix} \rho_2^2 & 0 & \rho_2\rho_4 \\ 0 & \rho_2^2 & 0 \\ \rho_2\rho_4 & 0 & \rho_4^2 + e^{2i\delta}\rho_3^2 \end{pmatrix}, \tag{16}$$

where

$$\rho_2 = \frac{m_2^\nu}{\sqrt{M_1}}, \quad \rho_4 = \frac{m_4^\nu}{\sqrt{M_1}}, \quad \rho_3 = \frac{m_3^\nu}{\sqrt{M_3}}. \tag{17}$$

The mass eigenvalues and diagonalization matrix of charged leptons are given by

$$V_l^\dagger M_l^* M_l^t V_l = diag(m_e^2, m_\mu^2, m_\tau^2) = ((m_2^l)^2, (m_3^l)^2, (m_1^l)^2), \tag{18}$$

$$V_l = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 & -1 & 1 \\ 0 & 1 & 1 \\ -\sqrt{2} & 0 & 0 \end{pmatrix}, \tag{19}$$

and those of neutrinos are given by

$$V_\nu^t M_\nu V_\nu = diag(e^{i(\phi_1-\phi)}m_{\nu_1}, e^{i(\phi_2+\phi)}m_{\nu_2}, m_{\nu_3}), \tag{20}$$

$$V_\nu = \begin{pmatrix} -\sin\theta_\nu & e^{i\phi}\cos\theta_\nu & 0 \\ 0 & 0 & 1 \\ e^{-i\phi}\cos\theta_\nu & \sin\theta_\nu & 0 \end{pmatrix}. \tag{21}$$

From Eq.(19) and Eq.(21), we obtain the MNS matrix as follows

$$V_{MNS} = V_l^\dagger V_\nu P_\nu = \frac{1}{\sqrt{2}}\begin{pmatrix} -\sqrt{2}e^{-i\phi}\cos\theta_\nu & -\sqrt{2}\sin\theta_\nu & 0 \\ \sin\theta_\nu & -e^{i\phi}\cos\theta_\nu & 1 \\ -\sin\theta_\nu & e^{i\phi}\cos\theta_\nu & 1 \end{pmatrix} P_\nu, \tag{22}$$

where

$$P_\nu = \mathrm{diag}(e^{-i(\phi_1-\phi)/2}, e^{-i(\phi_2+\phi)/2}, 1). \tag{23}$$

Here it is worth mentioning that the lower bound of $(0.04 <)\theta_{13}$ was shown by the recent experiment reported by T2K [15] at 90 % C.L., which could give a severe test to our model near future.

From the experimental bound [18], we impose the condition

$$\tan\theta_\nu = \frac{1}{\sqrt{2}}, \quad m_{\nu_2}^2 - m_{\nu_1}^2 = 8.0\times10^{-5}(\mathrm{eV}^2), \quad m_{\nu_2}^2 - m_{\nu_3}^2 = 2.5\times10^{-3}(\mathrm{eV}^2), \tag{24}$$

on the parameters, then the phase ϕ is given by

$$r\cos\phi = 0.361, \quad r = \frac{\rho_2}{\rho_4}. \tag{25}$$

Fixing the VEVs as

$$v_u = 10, \quad v_u' = 155.3, \quad v_d = 2.0, \quad v_d' = 77.8 \quad (\mathrm{GeV}), \tag{26}$$

and the charged lepton masses as [8][19]

$$m_1^l = 1.75\mathrm{GeV}, \quad m_2^l = 487\mathrm{keV}, \quad m_3^l = 103\mathrm{MeV}, \tag{27}$$

Yukawa coupling constants are given by

$$Y_1^E = 0.875, \quad Y_3^E = 5.15\times10^{-2}, \quad Y_2^E = 6.25\times10^{-6}. \tag{28}$$

For the RHN mass parameters, we assume

$$V = 10^{12}\mathrm{GeV}, \quad Y_1^M = Y_3^M = 1. \tag{29}$$

In order to investigate model dependence, we give two sample parameter sets A and B which are defined as follows

$$\begin{array}{lll} \mathrm{A} & : & r = 0.361 \\ & & \phi = \phi_1 = \phi_2 = 0.0^\circ, \quad \delta = 90.0^\circ, \\ & & \rho_2^2 = 1.80\times10^{-2}\mathrm{eV}, \quad \rho_3^2 = 14.47\times10^{-2}\mathrm{eV}, \quad \rho_4^2 = 13.78\times10^{-2}\mathrm{eV}, \\ & & m_{\nu_1} = 5.24\times10^{-2}\mathrm{eV}, \quad m_{\nu_2} = 5.31\times10^{-2}\mathrm{eV}, \quad m_{\nu_3} = 1.80\times10^{-2}\mathrm{eV}, \\ & & m_2^\nu = 4.24\mathrm{GeV}, \quad m_3^\nu = 12.0\mathrm{GeV}, \quad m_4^\nu = 11.7\mathrm{GeV}, \\ & & Y_2^N = 0.424, \quad Y_3^N = 0.077, \quad Y_4^N = 1.17, \end{array} \tag{30}$$

$$\begin{array}{lll} \mathrm{B} & : & r = 1.000 \\ & & \phi = 68.84^\circ, \quad \phi_1 = 41.55^\circ, \quad \phi_2 = 41.15^\circ, \quad \delta = 89.805^\circ, \\ & & \rho_2^2 = 5.03\times10^{-2}\mathrm{eV}, \quad \rho_3^2 = 10.04\times10^{-2}\mathrm{eV}, \quad \rho_4^2 = 5.03\times10^{-2}\mathrm{eV}, \\ & & m_{\nu_1} = 7.08\times10^{-2}\mathrm{eV}, \quad m_{\nu_2} = 7.14\times10^{-2}\mathrm{eV}, \quad m_{\nu_3} = 5.03\times10^{-2}\mathrm{eV} \\ & & m_2^\nu = 7.09\mathrm{GeV}, \quad m_3^\nu = 10.02\mathrm{GeV}, \quad m_4^\nu = 7.09\mathrm{GeV}, \\ & & Y_2^N = 0.709, \quad Y_3^N = 0.065, \quad Y_4^N = 0.709. \end{array} \tag{31}$$

Generally, multi-Higgs model causes FCNC problems, however our assignments do not cause such problems. In the lepton sector, the interactions between charged leptons and Higgs bosons are given by

$$\begin{aligned} \mathcal{L}_l &= Y_1^E \tau^c \left[l_\mu \left(\frac{H_2^D - H_1^D}{\sqrt{2}} \right) + l_\tau \left(\frac{H_1^D + H_2^D}{\sqrt{2}} \right) \right] - Y_2^E H_3^D e^c l_e \\ &+ Y_3^E \mu^c \left[l_\mu \left(\frac{H_1^D + H_2^D}{\sqrt{2}} \right) + l_\tau \left(\frac{H_1^D - H_2^D}{\sqrt{2}} \right) \right], \end{aligned} \tag{32}$$

which do not contribute to $\tau \to e + \gamma, \mu \to e + \gamma$ processes. Because $\mathcal{L}_l$ has accidental S_2 symmetry such as

$$(H_1^D, H_2^D) \to (H_2^D, H_1^D), \quad (l_\mu, \mu^c) \to (-l_\mu, -\mu^c), \tag{33}$$

$\tau \to \mu + \gamma$ process is also not induced. Note that this S_2 symmetry is not the symmetry of whole theory, as the symmetry is violated in neutrino Yukawa couplings and flavon superpotential W_Φ.

In the quark sector, as the quarks couple only to H_3^U, H_3^D, Higgs mediated FCNCs are not induced. In the basis that quark mass matrices are diagonal, the superpotential is written as

$$\begin{aligned} W_Q &= Y_t H_3^U (Q_3)'(U_3^c)' + Y_c H_3^U (Q_2)'(U_2^c)' + Y_u H_3^U (Q_1)'(U_1^c)' \\ &+ Y_b H_3^D (Q_3)'(D_3^c)' + Y_s H_3^D (Q_2)'(D_2^c)' + Y_d H_3^D (Q_1)'(D_1^c)'. \end{aligned} \tag{34}$$

From here, we fix top, bottom and charm masses as [8][19]

$$Y_t v_u' = 172.5, \quad Y_b v_d' = 2.89, \quad Y_c v_u' = 0.624 \quad (\mathrm{GeV}), \tag{35}$$

and assume exotic quark mass as

$$kv'_s = 2000 \quad (\mathrm{GeV}), \tag{36}$$

in order to forbid the decay of lightest flavon into exotic quark pair. Then we fix the values of Yukawa coupling constants as

$$Y_t = 1.12, \quad Y_b = 0.0371, \quad Y_c = 0.00405, \quad k = 1.0, \quad v'_s = 2000\mathrm{GeV}, \quad v_s = 200\mathrm{GeV}. \tag{37}$$

2.4 Higgs sector

Higgs potential is given as follows,

$$\begin{aligned}
V &= V_F + V_D + V_A + V_{m^2}, && (38)\\
V_F &= \left|\lambda_1 S_3 H_1^D + \lambda_5 S_1 H_3^D\right|^2 + \left|\lambda_1 S_3 H_2^D + \lambda_5 S_2 H_3^D\right|^2 \\
&+ \left|\lambda_3 S_3 H_3^D + \lambda_4 (S_1 H_1^D + S_2 H_2^D)\right|^2 \\
&+ \left|\lambda_1 S_3 H_1^U + \lambda_4 S_1 H_3^U\right|^2 + \left|\lambda_1 S_3 H_2^U + \lambda_4 S_2 H_3^U\right|^2 \\
&+ \left|\lambda_3 S_3 H_3^U + \lambda_5 (S_1 H_1^U + S_2 H_2^U)\right|^2 \\
&+ \left|\lambda_4 H_3^U H_1^D + \lambda_5 H_1^U H_3^D\right|^2 + \left|\lambda_4 H_3^U H_2^D + \lambda_5 H_2^U H_3^D\right|^2 \\
&+ \left|\lambda_1 (H_1^U H_1^D + H_2^U H_2^D) + \lambda_3 H_3^U H_3^D\right|^2, && (39)\\
V_D &= \frac{1}{2} g_Y^2 \left(\frac{1}{2}|H_a^U|^2 - \frac{1}{2}|H_a^D|^2\right)^2 + \frac{1}{2} g_2^2 \sum_A \left((H_a^U)^\dagger T^A H_a^U + (H_a^D)^\dagger T^A H_a^D\right)^2 \\
&+ \frac{1}{2} g_x^2 \left(-2|H_a^U|^2 - 3|H_a^D|^2 + 5|S_a|^2\right)^2, && (40)\\
V_A &= -\lambda_1 A_1 S_3 (H_1^U H_1^D + H_2^U H_2^D) - \lambda_3 A_3 S_3 H_3^U H_3^D \\
&- \lambda_4 A_4 H_3^U (S_1 H_1^D + S_2 H_2^D) - \lambda_5 A_5 (S_1 H_1^U + S_2 H_2^U) H_3^D + h.c., && (41)\\
V_{m^2} &= -m_{U3}^2 |H_3^U|^2 + m_U^2 (|H_1^U|^2 + |H_2^U|^2) + m_{D3}^2 |H_3^D|^2 + m_D^2 (|H_1^D|^2 + |H_2^D|^2) \\
&- m_{S3}^2 |S_3|^2 + m_S^2 (|S_1|^2 + |S_2|^2) \\
&- \left[m_{BU}^2 (H_3^U)^\dagger (H_1^U + H_2^U) + m_{BD}^2 (H_3^D)^\dagger (H_1^D + H_2^D) + m_{BS}^2 (S_3)^\dagger (S_1 + S_2) + h.c.\right], && (42)
\end{aligned}$$

where we can define $\lambda_{1,3,4,5}$ to be real without loss of generality, and we assume all the soft SUSY breaking parameters are real to avoid complex VEVs.

he soft $S_4 \times Z_2$ breaking terms; $m_{BU}^2, m_{BD}^2, m_{BS}^2$, violate accidental $O(2)$ symmetry of Higgs potential and fix the VEV directions (Eq.(13)) to realize $\theta_{23} = 45°$. This potential has S_2 symmetry such as

$$H_1^U \leftrightarrow H_2^U, \quad H_1^D \leftrightarrow H_2^D, \quad S_1 \leftrightarrow S_2. \tag{43}$$

Minimizing this potential, we get mass matrices of Higgs bosons. The results are given in appendix A. In the same manner, we add soft $S_4 \times Z_2$ breaking terms in flavon sector to avoid domain wall problem [20][1].

3 Dark Matter

Here we show that our model is consistent with cosmic-ray observation of PAMELA. Decaying dark matter scenarios with Non-Abelian discrete flavor symmetries have been done by Ref. [16].

3.1 LF decay width

We assume that the candidate for decaying dark matter is the lightest flavon (LF). In the six flavon superfields; $\Phi_a, \Phi_a^c (a = 1, 2, 3)$, only one linear combination is super-heavy and the other five superfields have TeV scale masses. As LF cannot decay into other flavons, it has very long lifetime. Due to the non-renormalizable interactions with light particles, LF becomes unstable dark matter. Among the interactions, the source term of RHN mass

---

[1] We would like to thank Refree for the suggestion.

$$W_{\text{eff}} = \frac{(Y^N H^U L)^2}{2Y^M(V+\Phi_3)} \sim \frac{(Y^N H^U L)^2}{2M_R}\left(1-\frac{\Phi_3}{V}\right) = \frac{1}{2}m_\nu\left(\frac{H^U L}{v}\right)^2\left(1-\frac{\Phi_3}{V}\right) \tag{44}$$

is the unique interaction to emit leptons without emitting quarks [21], where v is VEV of H^U. We estimate the positron flux using this interaction. Due to the factor $1/v$, the Higgs which develops the smallest VEV gives the largest contribution to LF decay. Therefore we can neglect the contribution from H_3^U, because $v_u \ll v_u'$ as one can see from Eq.(26). This effect is impotant to suppress weak boson emission. Due to the enhancement factor m_{LF}/v_u, LF decay width is dominated by 4-body decay as follows

$$\begin{align}
\Gamma(LF \to \nu+\nu) &\ll \Gamma(LF \to \nu+l+H^+) \sim \Gamma(LF \to \nu+l+W^+) \nonumber \\
&\ll \Gamma(LF \to l+l+H^+ +H^+). \tag{45}
\end{align}$$

From the spectrum of positron flux observed by PAMELA, we assume

$$m_{LF} = 4\text{TeV}. \tag{46}$$

If we assume all sfermions which couple to LF are heavier than 4TeV, the other interactions do not contribute to LF decay. The interactions which contribute to LF decay is given as follows

$$\begin{align}
\mathcal{L}_{2\nu} &= \frac{1}{2}C_{LF}\phi_{LF}\left\{[(H_1^U)^0(H_1^U)^0+(H_2^U)^0(H_2^U)^0](\nu_e\nu_e+r^2\nu_\mu\nu_\mu+r^2\nu_\tau\nu_\tau)\right. \nonumber \\
&+ \left. 2\sqrt{2}r(\nu_\mu-\nu_\tau)(H_1^U)^0(H_2^U)^0\nu_e-\sqrt{2}r(\nu_\mu+\nu_\tau)[(H_1^U)^0(H_1^U)^0-(H_2^U)^0(H_2^U)^0]\nu_e\right\}, \tag{47} \\
\mathcal{L}_{l\nu} &= -\frac{1}{2}C_{LF}\phi_{LF}\left\{2[(H_1^U)^0(H_1^U)^+ +(H_2^U)^0(H_2^U)^+](e\nu_e+r^2\nu_\mu\mu+r^2\nu_\tau\tau)\right. \nonumber \\
&+ 2r^2[(H_1^U)^0(H_2^U)^+ -(H_2^U)^0(H_1^U)^+](\nu_\mu\tau-\nu_\tau\mu) \nonumber \\
&+ \sqrt{2}r[(H_1^U)^0(H_2^U)^+ +(H_2^U)^0(H_1^U)^+][(\nu_\mu-\nu_\tau)e+\nu_e(\mu-\tau)] \nonumber \\
&- \left.\sqrt{2}r[(H_1^U)^0(H_1^U)^+ -(H_2^U)^0(H_2^U)^+][(\nu_\mu+\nu_\tau)e+\nu_e(\mu+\tau)]\right\}, \tag{48} \\
\mathcal{L}_{2l} &= \frac{1}{2}C_{LF}\phi_{LF}\left\{[(H_1^U)^+(H_1^U)^+ +(H_2^U)^+(H_2^U)^+](ee+r^2\mu\mu+r^2\tau\tau)\right. \nonumber \\
&+ \left. 2\sqrt{2}r(\mu-\tau)(H_1^U)^+(H_2^U)^+e-\sqrt{2}r(\mu+\tau)[(H_1^U)^+(H_1^U)^+ -(H_2^U)^+(H_2^U)^+]e\right\}, \tag{49} \\
C_{LF} &= \frac{\epsilon\rho_4^2}{\sqrt{2}Vv_u^2}, \tag{50}
\end{align}$$

where ϵ is the flavon mixing parameter which is defined by

$$\Phi_3 = \frac{\epsilon\phi_{LF}}{\sqrt{2}}, \tag{51}$$

where ϕ_{LF} is LF field.

Using Eq.(47)-(49), the LF decay widths are given as follows

$$\begin{align}
\Gamma_{2\nu} &= \Gamma_{2\bar{\nu}} = (6+10r^2+12r^4)\Gamma_0, \tag{52} \\
\Gamma_{l\nu} &= \Gamma_{\bar{l}\bar{\nu}} = \Gamma_e+\Gamma_\mu+\Gamma_\tau = (2+8r^2+8r^4)\Gamma_0, \nonumber \\
&\quad \Gamma_e = \Gamma_{\bar{e}} = (2+4r^2)\Gamma_0, \quad \Gamma_\mu = \Gamma_{\bar{\mu}} = \Gamma_\tau = \Gamma_{\bar{\tau}} = (2r^2+4r^4)\Gamma_0, \nonumber \\
\Gamma_{2l} &= \Gamma_{2\bar{l}} = \Gamma_{2e}+\Gamma_{2\mu}+\Gamma_{2\tau}+\Gamma_{e\mu}+\Gamma_{e\tau} = (8+12r^2+16r^4)\Gamma_0, \tag{53} \\
&\quad \Gamma_{2e} = 8\Gamma_0, \quad \Gamma_{2\mu} = \Gamma_{2\tau} = 8r^4\Gamma_0, \quad \Gamma_{e\mu} = \Gamma_{e\tau} = 6r^2\Gamma_0, \tag{54} \\
\Gamma_{\text{lepton}} &= \Gamma_{l\nu}+\Gamma_{2l} = (10+20r^2+24r^4)\Gamma_0 = \Gamma_{\text{anti-lepton}}, \nonumber \\
\Gamma_{\text{total}} &= 2(\Gamma_{2\nu}+\Gamma_{l\nu}+\Gamma_{2l}) = 2(16+30r^2+36r^4)\Gamma_0, \tag{55} \\
\Gamma_0 &= \frac{m_{LF}}{16\pi}\left(\frac{m_{LF}^2\epsilon\rho_4^2}{32\pi^2 v_u^2 V}\right)^2(0.111), \tag{56}
\end{align}$$

where we classify the final states only by charged lepton flavor e, μ, τ. The rates of lepton flavor emitted by LF decay are given by

$$p_e = \frac{9+8r^2}{9+16r^2+20r^4}, \quad p_\mu = p_\tau = \frac{4r^2+10r^4}{9+16r^2+20r^4}. \tag{57}$$

Anti-lepton flux depends on Majorana phase ϕ through Eq.(25), such as e-dominant for $r < 1$ and (μ, τ)-dominant for $r > 1$ (see Fig. 1). For each parameter set, LF lifetime is estimated as follows

$$\mathrm{A} \;:\; \Gamma^{-1}_{\text{total}} = 3.72 \times 10^{11} \epsilon^{-2} \text{sec}, \quad \Gamma^{-1}_{\text{anti-lepton}} = 1.17 \times 10^{12} \epsilon^{-2} \text{sec}, \tag{58}$$

$$\mathrm{B} \;:\; \Gamma^{-1}_{\text{total}} = 7.01 \times 10^{11} \epsilon^{-2} \text{sec}, \quad \Gamma^{-1}_{\text{anti-lepton}} = 2.13 \times 10^{12} \epsilon^{-2} \text{sec}. \tag{59}$$

Hereafter we assume $\epsilon < 10^{-3}$ to avoid extinction of LF.



Figure 1:

3.2 Relic Abundance of LF

At early stage of the universe, flavon multiplets are produced through $U(1)_Z$ gauge interaction [13]. Since we assume that reheating temperature is low enough to avoid gravitino over-production as $T_{RH} < 10^7\text{GeV}$ [22], this interaction is never thermal equiliburium. Therefore we assume non-thermal production of flavons and boundary condition $n_{LF}(T_{RH}) = 0$.

For the chiral multiplets (ψ_L, Ψ), $U(1)_Z$ gauge interaction is given by

$$\mathcal{L}_{U(1)_Z} = i g_z A^\mu \sum_i z_i \left[\bar{\psi}_{i,L} \gamma_\mu \psi_{i,L} + \Psi_i \partial_\mu \Psi_i^\dagger - \Psi_i^\dagger \partial_\mu \Psi_i \right], \tag{60}$$

from which we calculate production cross sections of flavon multiplets (ϕ, Φ). From Eq.(13), the $U(1)_Z$ gauge boson mass is nearly equal to $16 g_z V$.

As all produced flavon multiplets decay into LF finally, LF number density is given by

$$n_{LF} = 5 N_{LF} (n_\phi + n_\Phi), \tag{61}$$

where 5 is the number of light flavon superfields, n_Φ, n_ϕ are number density of one flavon multiplet and $N_{LF} \sim O(1)$ is LF production rate which means how many LFs are produced per one degree of freedom of flavon multiplets. The Boltzmann equation for n_{LF} is given by

$$\begin{aligned} \dot{n}_{LF} + 3 H n_{LF} &= 2480 N_{LF} C T^8, & (62) \\ C &= \frac{21}{(2\pi)^5} \left(\frac{1}{32 V^2} \right)^2, & (63) \end{aligned}$$

from which we get

$$\begin{aligned} \Omega_{LF} h^2 &= \frac{m_{LF} s_0 h^2}{\rho_c} \left[\frac{15 \times 2480 \times 21 m_P N_{LF}}{2\pi^2 (341.25) \times 30.67 (2\pi)^5} \left(\frac{1}{32 V^2} \right)^2 T_{RH}^3 \right] \\ &= 5.06 \times 10^{-9} N_{LF} \left(\frac{T_{RH}}{10^5 GeV} \right)^3, & (64) \end{aligned}$$

where [18]

$$
\begin{aligned}
H &= 1.66\sqrt{g_*}\frac{T^2}{m_P}, \\
g_* &= 341.25, \\
m_P &= 1.22 \times 10^{19}\mathrm{GeV}, \\
s_0 &= 2890/\mathrm{cm}^3, \\
\rho_c &= 1.05 \times 10^4 h^2 \mathrm{eV}/\mathrm{cm}^3. \qquad (65)
\end{aligned}
$$

For $T_{RH} < 10^7$GeV, LF does not dominate dark matter ($\Omega_{LF}h^2 \ll \Omega_{DM}h^2 = 0.11$), thus other dark matter should be considered as we will discuss later. Such multi-component dark matter is discussed in [26]. Although the number density is very low, the short lifetime of LF enables us to explain cosmic-ray observation. The effective lifetime of LF is defined as

$$\tau_{\mathrm{eff}} \equiv \Gamma^{-1}_{\text{anti-lepton}}\left(\frac{\Omega_{DM}}{\Omega_{LF}}\right), \tag{66}$$

and the following values are obtained for each parameter set

$$\mathrm{A} \quad : \quad \tau_{\mathrm{eff}} = 6.0 \times 10^{25}\mathrm{sec}, \quad \epsilon^2 N_{LF}\left(\frac{T_{RH}}{10^5\mathrm{GeV}}\right)^3 = 4.2 \times 10^{-7}, \tag{67}$$

$$\mathrm{B} \quad : \quad \tau_{\mathrm{eff}} = 7.0 \times 10^{25}\mathrm{sec}, \quad \epsilon^2 N_{LF}\left(\frac{T_{RH}}{10^5\mathrm{GeV}}\right)^3 = 6.6 \times 10^{-7}. \tag{68}$$

Eq.(67) and Eq.(68) are satisfied for example, if we put $N_{LF} \sim 1, T_{RH} \sim 10^5\mathrm{GeV}, \epsilon \sim 10^{-3}$.

The positron flux from the decay of LF is calculated as

$$\Phi(E_{e^+}) = \frac{v_{e^+}}{4\pi}\frac{1}{m_{LF}\tau_{\mathrm{eff}}}\int dE' G_{e^+}(E_{e^+}, E') \sum_{\ell=e^+,\mu^+,\tau^\pm} p_\ell \frac{dN_{\ell\ e^+}}{dE'}, \tag{69}$$

where v_{e^+} is the velocity of the positron, G_{e^+} is the Green's fuction which is expressed in [3], p_ℓ is expressed in Eq.(57) and $dN_{\ell\ e^+}/dE'$ is the fragmentation function produced from the decay of ℓ to e^+. The fragmentation function is calculated by using the event generator `pythia` [27] and the result is shown in Fig.2. We can evalutate the positron flux from the decay of LF by using the fragmentation function. The results for each parameter set A and B are shown in Fig.3.



Figure 2: The fragmentation function calculated by `pythia` for parameter set A (left) and B (right).

From the gamma-ray observations [4], the constraint for τ-flux is given by

$$(\tau_\tau)_{\mathrm{eff}} = \frac{\Gamma_{\mathrm{lepton}}}{\Gamma_\tau}\tau_{\mathrm{eff}} = \frac{10 + 20r^2 + 24r^4}{8r^2 + 20r^4}\tau_{\mathrm{eff}} \geq 2.1 \times 10^{26}\mathrm{sec}, \tag{70}$$



Figure 3: The positron flux calculated for parameter set A (left) and B (right).

which is estimated for each parameter set as follows

$$\begin{aligned} \mathrm{A} \quad &: \quad (\tau_\tau)_{\rm eff} = 5.6 \times 10^{26}\mathrm{sec}, \\ \mathrm{B} \quad &: \quad (\tau_\tau)_{\rm eff} = 1.4 \times 10^{26}\mathrm{sec}. \end{aligned} \tag{71}$$

As the parameter set B is severe to satisfy Eq.(70), small r model is favored. This is the new information about neurtino sector extarcted by cosmic-ray observations.

3.3 Higgs decay width

No excess of anti-proton flux in cosmic-ray constrains the species of the particles emitted by LF decay [23]. As the weak boson $Z, W^\pm$ and the chargino decay mainly into quarks, LF should not decay into these particles so much. The weak boson emission is suppressed by factor $(v_u/m_{LF})^2$ and the chargino emission channel is kinematically closed for heavy sfermion scenario. In order to forbid the weak boson and the chargino emission from Higgs boson decay, we assume light Higgs scenario.

In the Higgs potential Eqs.(38)-(42) and mass terms of the neutralinos and the charginos

$$\begin{aligned} \mathcal{L} \quad \supset \quad & -i\sqrt{2}(H_a^U)^\dagger[g_2\lambda_2^A T_2^A + 3g_y\lambda_Y - 2g_x\lambda_X]h_a^U \\ - \quad & i\sqrt{2}(H_a^D)^\dagger[g_2\lambda_2^A T_2^A - 3g_y\lambda_Y - 3g_x\lambda_X]h_a^D - i\sqrt{2}(S_a)^\dagger[5g_x\lambda_X]s_a \\ - \quad & \frac{1}{2}M_2\lambda_2^A\lambda_2^A - \frac{1}{2}M_Y\lambda_Y\lambda_Y - \frac{1}{2}M_X\lambda_X\lambda_X + (W_H)_{\theta^2} + h.c. \quad (A = 1, 2, 3), \end{aligned} \tag{72}$$

we assume the parameters as follows

$$\begin{aligned} & g_2 = 0.652, \quad g_y = \frac{1}{6}g_Y \quad (g_Y = 0.357), \quad g_x = \frac{1}{2\sqrt{6}}g_Y = 0.073, \\ & \lambda_1 = 0.065, \quad \lambda_3 = 0.4, \quad \lambda_4 = 0.398, \quad \lambda_5 = 0.75, \\ & M_X = M_2 = 200, \quad M_Y = 180 \quad (\mathrm{GeV}), \\ & A_1 = 0.0, \quad A_3 = A_4 = A_5 = 1.0 \quad (\mathrm{TeV}), \\ & m_{BU}^2 = 0.0408, \quad m_{BD}^2 = m_{BS}^2 = 0.02 \quad (\mathrm{TeV}^2). \end{aligned} \tag{73}$$

Mass matrices of neutralinos and charginos are given in appendix A and the values of mass eigenvalues and mixing matrices are given in appendix B.

We consider only the mass eigenstates which dominate $H_{1,2}^U$ such as

$$\begin{aligned} & \phi_1'(91.50), \quad \phi_4'(121.96), \quad \phi_5'(152.48), \quad \rho_1'(112.12), \quad \rho_6'(145.53), \\ & (H_1^\pm)'(90.70), \quad (H_3^\pm)'(130.02) \quad (\mathrm{GeV}), \end{aligned} \tag{74}$$

where $\phi_1', \rho_1', (H_1^\pm)'$ are S_2-odd and the others are even. As S_2 forbids interaction $\phi_1' ZZ$ and ϕ_1' is not emitted through $Z^* \to Z + \phi_1'$, LEP bound $m_H \geq 114.4$GeV is not imposed on ϕ_1'. As the masses of these Higgs bosons are well degenerated, they do not emit weak bosons or charginos.

The neutralinos into which these Higgs bosons can decay are two singlino dominant neutralinos

$$\eta_1'(41.92), \quad \eta_4'(44.55) \quad (\mathrm{GeV}), \tag{75}$$

where η_1' is S_2-odd and LSP. S_2-even neutralino η_4' can decay into η_1' through $\eta_4' \to \eta_1' + \mu + \bar{\tau}$ without emitting quarks. As S_2-odd Higgs boson can not decay into quarks, we consider only the decay of S_2-even Higgs bosons.

The decay widths of $\phi_4', \phi_5', \rho_6'$ due to the Yukawa interactions

$$\mathcal{L} \supset Y_c(H_3^U)^0 cc^c + Y_b(H_3^D)^0 bb^c + Y_1^E[(H_1^D)^0 e_1^c e_1 + (H_2^D)^0 e_1^c e_2] + h.c., \tag{76}$$

are given as follows

$$\begin{aligned}
\Gamma(\phi_4' \to c + \bar{c}) &= 3.63 \times 10^{-6}\Gamma_{2d}, && (77)\\
\Gamma(\phi_4' \to b + \bar{b}) &= 1.54 \times 10^{-4}\Gamma_{2d}, && (78)\\
\Gamma(\phi_4' \to \tau + \bar{\tau}) &= 1.10 \times 10^{-6}\Gamma_{2d}, && (79)\\
\Gamma(\phi_5' \to c + \bar{c}) &= 1.61 \times 10^{-5}\Gamma_{2d}, && (80)\\
\Gamma(\phi_5' \to b + \bar{b}) &= 2.68 \times 10^{-4}\Gamma_{2d} && (81)\\
\Gamma(\phi_5' \to \tau + \bar{\tau}) &= 4.90 \times 10^{-5}\Gamma_{2d}, && (82)\\
\Gamma(\rho_6' \to c + \bar{c}) &= 1.91 \times 10^{-7}\Gamma_{2d}, && (83)\\
\Gamma(\rho_6' \to b + \bar{b}) &= 4.96 \times 10^{-6}\Gamma_{2d}, && (84)\\
\Gamma(\rho_6' \to \tau + \bar{\tau}) &= 6.89 \times 10^{-8}\Gamma_{2d}, && (85)
\end{aligned}$$

where Γ_{2d} is 2-body decay width of scalar. The interactions with the neutralinos η_1', η_4',

$$\begin{aligned}
\mathcal{L} \supset\ & -\lambda_1 \left\{S_3[(h_1^U)^0(h_1^D)^0 + (h_2^U)^0(h_2^D)^0] + s_3[(H_1^U)^0(h_1^D)^0 + (H_2^U)^0(h_2^D)^0]\right.\\
+\ & \left. s_3[(h_1^U)^0(H_1^D)^0 + (h_2^U)^0(H_2^D)^0]\right\}\\
-\ & \lambda_3[S_3(h_3^U)^0(h_3^D)^0 + s_3(H_3^U)^0(h_3^D)^0 + s_3(h_3^U)^0(H_3^D)^0]\\
-\ & \lambda_4 \left\{(H_3^U)^0[s_1(h_1^D)^0 + s_2(h_2^D)^0] + (h_3^U)^0[S_1(h_1^D)^0 + S_2(h_2^D)^0]\right.\\
+\ & \left. (h_3^U)^0[s_1(H_1^D)^0 + s_2(H_2^D)^0]\right\}\\
-\ & \lambda_5 \left\{(H_3^D)^0[s_1(h_1^U)^0 + s_2(h_2^U)^0] + (h_3^D)^0[S_1(h_1^U)^0 + S_2(h_2^U)^0]\right.\\
+\ & \left. (h_3^D)^0[s_1(H_1^U)^0 + s_2(H_2^U)^0]\right\}\\
-\ & i\sqrt{2}\sum_i [(H_i^U)^0]^\dagger \left[-\frac{1}{2}g_2\lambda_2^3 + \frac{1}{2}g_Y\lambda_Y - 2g_x\lambda_X\right](h_i^U)^0\\
-\ & i\sqrt{2}\sum_i [(H_i^D)^0]^\dagger \left[\frac{1}{2}g_2\lambda_2^3 - \frac{1}{2}g_Y\lambda_Y - 3g_x\lambda_X\right](h_i^D)^0\\
-\ & i\sqrt{2}\sum_i S_i^\dagger[5g_x\lambda_X]s_i + h.c.\\
=\ & -(0.0620\phi_4' + 0.127\phi_5' + 0.0299i\rho_6')\eta_4'\eta_4'\\
-\ & (-0.0716\phi_4' + 0.117\phi_5' + 0.0142i\rho_6')\eta_1'\eta_1' + h.c.
\end{aligned} \tag{86}$$

give

$$\begin{aligned}
\Gamma(\phi_4' \to \eta + \eta) &= 208 \times 10^{-4}\Gamma_{2d}, && (87)\\
\Gamma(\phi_5' \to \eta + \eta) &= 864 \times 10^{-4}\Gamma_{2d}, && (88)\\
\Gamma(\rho_6' \to \eta + \eta) &= 3179 \times 10^{-6}\Gamma_{2d}, && (89)
\end{aligned}$$

which dominate the decay widths of $\phi_4', \phi_5', \rho_6'$.

The decay widths of $(H_3^\pm)'$ due to Yukawa interactions

$$\mathcal{L} \supset -Y_c(H_3^U)^+ sc^c - Y_1^E[(H_1^D)^- e^c \nu_1 + (H_2^D)^- e_1^c \nu_2] + h.c. \tag{90}$$

are given by

$$\begin{aligned}\Gamma((H_3^-)' \to s + \bar{c}) &= 1.95 \times 10^{-7}\Gamma_{2d}, &(91)\\ \Gamma((H_3^-)' \to \tau + \bar{\nu}_\tau) &= 2.48 \times 10^{-6}\Gamma_{2d}. &(92)\end{aligned}$$

From these estimations, $(H_3^\pm)'$ decay gives dominant contribution to anti-proton flux.

As one charged lepton emission from LF decay accommodates one charged Higgs emission at even rate of $(H_1^\pm)'$ and $(H_3^\pm)'$, the quark flux is estimated as

$$\tau_{\rm quark} = \left[\frac{1}{2}\left(0 + \frac{0.195}{2.48 + 0.195}\right)\right]^{-1} \tau_{\rm eff} = 27.4\tau_{\rm eff}. \quad (93)$$

For each parameter sets, we get

$$\begin{aligned}\mathrm{A} &: \tau_{\rm quark} = 1.6 \times 10^{27}\mathrm{sec}, &(94)\\ \mathrm{B} &: \tau_{\rm quark} = 1.9 \times 10^{27}\mathrm{sec}, &(95)\end{aligned}$$

from which the spectrum of anti-proton flux is given in Fig. 4. There is no inconsistency in anti-proton flux.



Figure 4:

3.4 Relic Abundance of LSP

Finally we estimate the relic abundance of LSP. The interactions between η_1', η_4' and Z are given by

$$\begin{aligned}\mathcal{L} \supset\ & \frac{1}{2}\bar{\psi}_1\gamma^\mu(-\partial_\mu - iG_1 Z_\mu\gamma_5)\psi_1 + \frac{1}{2}\bar{\psi}_4\gamma^\mu(-\partial_\mu - iG_4 Z_\mu\gamma_5)\psi_4\\ +\ & iG(f_L)\bar{\psi}_f\gamma^\mu Z_\mu P_L\psi_f + iG(f_R)\bar{\psi}_f\gamma^\mu Z_\mu P_R\psi_f, \quad (96)\end{aligned}$$

where

$$G_1 = 0.00823, \quad G_4 = 0.0119, \quad (97)$$

$$G(e_L) = -\frac{g_Y^2}{\sqrt{g_Y^2 + g_2^2}} = -0.172, \quad G(e_R) = \frac{-g_Y^2 + g_2^2}{2\sqrt{g_Y^2 + g_2^2}} = 0.200, \quad (98)$$

$$G(\nu_L) = -\frac{\sqrt{g_Y^2 + g_2^2}}{2} = -0.372, \quad (99)$$

$$G(u_L) = \frac{2g_Y^2}{3\sqrt{g_Y^2 + g_2^2}} = 0.114, \quad G(u_R) = \frac{g_Y^2 - 3g_2^2}{6\sqrt{g_Y^2 + g_2^2}} = -0.257, \quad (100)$$

$$G(d_L) = -\frac{g_Y^2}{3\sqrt{g_Y^2 + g_2^2}} = -0.057, \quad G(d_R) = \frac{g_Y^2 + 3g_2^2}{6\sqrt{g_Y^2 + g_2^2}} = 0.315. \quad (101)$$

These interactions give dominant contribution to annihilation of η'_1, η'_4. As $G_i \ll G(\nu_L)$, the contributions $Z \to \eta\eta$ to Z-decay width is negligible. Therefore LEP bound $m \geq 46$GeV is not imposed on η'_1, η'_4. The relic abundance of the neutralino is calculated by the formula

$$\begin{aligned} x_F &= \ln \frac{0.0955 m_P m_i (a + 6b/x_F)}{(g_* x_F)^{\frac{1}{2}}} \quad (i = 1, 4), && (102) \\ \Omega h^2 &= \frac{8.76 \times 10^{-11} g_*^{-\frac{1}{2}} x_F}{(a + 3b/x_F)\mathrm{GeV}^2}, && (103) \end{aligned}$$

where $m_1 = 41.92$GeV, $m_4 = 44.55$GeV,

$$\begin{aligned} a_{f,i} &= \frac{2c_f}{\pi}\left[\frac{m_f (G_i/2)}{4m_i^2 - M_Z^2}(G(f_L) - G(f_R))\right]^2 \left(1 - \frac{m_f^2}{m_i^2}\right)^{\frac{1}{2}}, && (104) \\ b_{f,i} &= \frac{1}{6}\left(-\frac{9}{2} + \frac{3}{4}\frac{m_f^2}{m_i^2 - m_f^2}\right) a_{f,i} \\ &+ \frac{c_f}{3\pi}\left[\frac{m_i (G_i/2)}{4m_i^2 - m_Z^2}\right]^2 \left[G^2(f_L) + G^2(f_R)\right]\left(4 + \frac{2m_f^2}{m_i^2}\right)\left(1 - \frac{m_f^2}{m_i^2}\right)^{\frac{1}{2}}, && (105) \end{aligned}$$

c_f is color factor such as $c_f = 1$ for $SU_c(3)$ singlet, $c_f = 3$ for triplet, and $m_Z = 91.2$ GeV. For the approximation $m_f/m_i = 0$, these coefficients are given by

$$\begin{aligned} a_1 &= a_{b,1} + a_{\tau,1} = 5.01 \times 10^{-11}\mathrm{GeV}^{-2}, && (106) \\ a_4 &= a_{b,4} + a_{\tau,4} = 1.22 \times 10^{-9}\mathrm{GeV}^{-2}, && (107) \\ b_1 &= 1.53 \times 10^{-8}\mathrm{GeV}^{-2}, && (108) \\ b_4 &= 4.23 \times 10^{-7}\mathrm{GeV}^{-2}. && (109) \end{aligned}$$

The relic abundance of η'_1 is given by

$$\begin{aligned} g_* &= 75.75, \\ x_F &= 22.32, \\ T_F &= 1.88 \text{ GeV}, \\ \Omega_1 h^2 &= 0.106, && (110) \end{aligned}$$

and that of η'_4 is given by

$$\begin{aligned} g_* &= 72.25, \\ x_F &= 25.52, \\ T_F &= 1.75 \text{ GeV}, \\ \Omega_4 h^2 &= 0.0052. && (111) \end{aligned}$$

As η'_4 is converted into η'_1, relic abundance of LSP is given by

$$(\Omega_{CDM} h^2) = \Omega_1 h^2 + \Omega_4 h^2 = 0.111, \qquad (112)$$

which realizes density parameter of dark matter.

4 Conclusion

We have considered dark matter based on $S_4 \times Z_2$ flavor symmetric extra U(1) model. The results are as follows. There exists appropriate parameter set to realize relic abundance of dark matter and positron flux observed by PAMELA at the same time. The dominant component of dark matter is LSP and the origin of positron flux is given by the decay of LF which generates the mass of RHN. There is deep connection between PAMELA observation and neutrino mass. The long life time of LF results in large RHN mass and the spectrum of positron flux depend on Majorana phase ϕ in V_{MNS}. Therefore, cosmic-ray observation gives new information about the structure of neutrino mass matrix.

From the fact that there is no excess of anti-proton flux in cosmic-ray, we can guess about the particle spectrum. As sfermions can decay into quarks, weak boson and charginos, LF must not decay into those particles, which sugests sfermions are heavier than 4TeV. Although this is also favorable from the viewpoint of the FCNC constraints, experimental verification becomes difficult. However experimantal verification of our scenario is not imposssible. From the fact that Higgs does not decay into weak boson or charginos, we can expect that Higgs boson is light and degenerated, therefore the examination of the mass spectrum of Higgs boson is possible.

A Mass matrices

Neutral CP even Higgs boson

$$\begin{aligned}
H^U_a &\supset \frac{\phi_{U,a}}{\sqrt{2}}, \quad H^D_a \supset \frac{\phi_{D,a}}{\sqrt{2}}, \quad S_a \supset \frac{\phi_{S,a}}{\sqrt{2}} \quad (a=1,2,3), \\
-\mathcal{L} &\supset \frac{1}{2}\phi_i M^2_{ij}\phi_j, \quad \phi_i = \begin{pmatrix} \phi_{U,a} \\ \phi_{D,a} \\ \phi_{S,a} \end{pmatrix}, \quad (i,j=1,2,\cdots,9),
\end{aligned}$$

$$\begin{aligned}
M^2_{1,1} &= M^2_{2,2} = \sqrt{2}m^2_{BU}(v'_u/v_u) + \lambda_1 A_1 v'_s(v_d/v_u) + \lambda_5 A_5 v_s(v'_d/v_u) - \lambda_1^2 v_d^2/2 - \lambda_5^2 v_s^2/2 \\
&- [(\lambda_1\lambda_4+\lambda_3\lambda_5)v_s v'_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v'_d v_d]\,(v'_u/v_u) + \left[\frac{1}{4}(g_Y^2+g_2^2)+4g_x^2\right] v_u^2, \\
M^2_{1,2} &= \lambda_5^2 v_s^2/2 + \lambda_1^2 v_d^2/2 + \left[\frac{1}{4}(g_Y^2+g_2^2)+4g_x^2\right] v_u^2, \\
M^2_{1,3} &= M^2_{2,3} = -m^2_{BU} + [(\lambda_1\lambda_4+\lambda_3\lambda_5)v'_s v_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v_d v'_d]/\sqrt{2} \\
&+ \sqrt{2}\left[\frac{1}{4}(g_Y^2+g_2^2)+4g_x^2\right] v_u v'_u, \\
M^2_{3,3} &= \sqrt{2}m^2_{BU}(v_u/v'_u) + \lambda_3 A_3 v'_s(v'_d/v'_u) + \lambda_4 A_4 v_s(v_d/v'_u) \\
&- [(\lambda_1\lambda_4+\lambda_3\lambda_5)v_s v'_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v'_d v_d]\,(v_u/v'_u) + 2\left[\frac{1}{4}(g_Y^2+g_2^2)+4g_x^2\right](v'_u)^2, \\
M^2_{4,4} &= M^2_{5,5} = \sqrt{2}m^2_{BD}(v'_d/v_d) + \lambda_1 A_1 v'_s(v_u/v_d) + \lambda_4 A_4 v_s(v'_u/v_d) - \lambda_1^2 v_u^2/2 - \lambda_4^2 v_s^2/2 \\
&- [(\lambda_1\lambda_5+\lambda_3\lambda_4)v_s v'_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v'_u v_u]\,(v'_d/v_d) + v_d^2\left[\frac{1}{4}(g_Y^2+g_2^2)+9g_x^2\right], \\
M^2_{4,5} &= \lambda_4^2 v_s^2/2 + \lambda_1^2 v_u^2/2 + \left[\frac{1}{4}(g_Y^2+g_2^2)+9g_x^2\right] v_d^2, \\
M^2_{4,6} &= M^2_{5,6} = -m^2_{BD} + [(\lambda_1\lambda_5+\lambda_3\lambda_4)v'_s v_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v_u v'_u]/\sqrt{2} \\
&+ \sqrt{2}\left[\frac{1}{4}(g_Y^2+g_2^2)+9g_x^2\right] v_d v'_d, \\
M^2_{6,6} &= \sqrt{2}m^2_{BD}(v_d/v'_d) + \lambda_3 A_3 v'_s(v'_u/v'_d) + \lambda_5 A_5 v_s(v_u/v'_d) \\
&- [(\lambda_1\lambda_5+\lambda_3\lambda_4)v_s v'_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v'_u v_u]\,(v_d/v'_d) + 2\left[\frac{1}{4}(g_Y^2+g_2^2)+9g_x^2\right](v'_d)^2, \\
M^2_{7,7} &= M^2_{8,8} = \sqrt{2}m^2_{BS}(v'_s/v_s) + \lambda_4 A_4(v'_u v_d/v_s) + \lambda_5 A_5(v_u v'_d/v_s) - \lambda_4^2 v_d^2/2 - \lambda_5^2 v_u^2/2 \\
&- [(\lambda_1\lambda_5+\lambda_3\lambda_4)v_d v'_d + (\lambda_1\lambda_4+\lambda_3\lambda_5)v'_u v_u]\,(v'_s/v_s) + \left[25g_x^2\right] v_s^2, \\
M^2_{7,8} &= \lambda_4^2 v_d^2/2 + \lambda_5^2 v_u^2/2 + \left[25g_x^2\right] v_s^2, \\
M^2_{7,9} &= M^2_{8,9} = -m^2_{BS} + [(\lambda_1\lambda_5+\lambda_3\lambda_4)v'_d v_d + (\lambda_1\lambda_4+\lambda_3\lambda_5)v_u v'_u]/\sqrt{2} + \sqrt{2}\left[25g_x^2\right] v_s v'_s, \\
M^2_{9,9} &= \sqrt{2}m^2_{BS}(v_s/v'_s) + \lambda_1 A_1(v_u v_d/v'_s) + \lambda_3 A_3(v'_u v'_d/v'_s) \\
&- [(\lambda_1\lambda_5+\lambda_3\lambda_4)v_d v'_d + (\lambda_1\lambda_4+\lambda_3\lambda_5)v'_u v_u]\,(v_s/v'_s) + 2\left[25g_x^2\right](v'_s)^2,
\end{aligned}$$

$$
\begin{aligned}
M^2_{1,4} &= M^2_{2,5} = -\lambda_1 A_1 v'_s + 3\lambda_1^2 v_u v_d/2 + (\lambda_4\lambda_5 + \lambda_1\lambda_3) v'_u v'_d + \left[-\frac{1}{4}(g_Y^2 + g_2^2) + 6g_x^2\right] v_u v_d, \\
M^2_{1,5} &= M^2_{2,4} = \lambda_1^2 v_u v_d/2 + \left[-\frac{1}{4}(g_Y^2 + g_2^2) + 6g_x^2\right] v_u v_d, \\
M^2_{1,6} &= M^2_{2,6} = -\lambda_5 A_5 v_s/\sqrt{2} + \sqrt{2}\lambda_5^2 v_u v'_d + (\lambda_4\lambda_5 + \lambda_1\lambda_3) v'_u v_d/\sqrt{2} \\
&+ \sqrt{2}\left[-\frac{1}{4}(g_Y^2 + g_2^2) + 6g_x^2\right] v_u v'_d, \\
M^2_{3,4} &= M^2_{3,5} = -\lambda_4 A_4 v_s/\sqrt{2} + \sqrt{2}\lambda_4^2 v'_u v_d + (\lambda_4\lambda_5 + \lambda_1\lambda_3) v_u v'_d/\sqrt{2} \\
&+ \sqrt{2}\left[-\frac{1}{4}(g_Y^2 + g_2^2) + 6g_x^2\right] v'_u v_d, \\
M^2_{3,6} &= -\lambda_3 A_3 v'_s + 2\lambda_3^2 v'_u v'_d + (\lambda_1\lambda_3 + \lambda_4\lambda_5) v_u v_d + 2\left[-\frac{1}{4}(g_Y^2 + g_2^2) + 6g_x^2\right] v'_u v'_d, \\
M^2_{1,7} &= M^2_{2,8} = -\lambda_5 A_5 v'_d + 3\lambda_5^2 v_u v_s/2 + (\lambda_1\lambda_4 + \lambda_3\lambda_5) v'_u v'_s + \left[-10g_x^2\right] v_u v_s, \\
M^2_{1,8} &= M^2_{2,7} = \lambda_5^2 v_u v_s/2 + \left[-10g_x^2\right] v_u v_s, \\
M^2_{1,9} &= M^2_{2,9} = -\lambda_1 A_1 v_d/\sqrt{2} + \sqrt{2}\lambda_1^2 v_u v'_s + (\lambda_1\lambda_4 + \lambda_3\lambda_5) v'_u v_s/\sqrt{2} + \sqrt{2}\left[-10g_x^2\right] v_u v'_s, \\
M^2_{3,7} &= M^2_{3,8} = -\lambda_4 A_4 v_d/\sqrt{2} + \sqrt{2}\lambda_4^2 v'_u v_s + (\lambda_1\lambda_4 + \lambda_3\lambda_5) v_u v'_s/\sqrt{2} + \sqrt{2}\left[-10g_x^2\right] v'_u v_s, \\
M^2_{3,9} &= -\lambda_3 A_3 v'_d + 2\lambda_3^2 v'_u v'_s + (\lambda_1\lambda_4 + \lambda_3\lambda_5) v_u v_s + 2\left[-10g_x^2\right] v'_u v'_s, \\
M^2_{4,7} &= M^2_{5,8} = -\lambda_4 A_4 v'_u + 3\lambda_4^2 v_d v_s/2 + (\lambda_1\lambda_5 + \lambda_3\lambda_4) v'_d v'_s + \left[-15g_x^2\right] v_d v_s, \\
M^2_{4,8} &= M^2_{5,7} = \lambda_4^2 v_d v_s/2 + \left[-15g_x^2\right] v_d v_s, \\
M^2_{4,9} &= M^2_{5,9} = -\lambda_1 A_1 v_u/\sqrt{2} + \sqrt{2}\lambda_1^2 v_d v'_s + (\lambda_1\lambda_5 + \lambda_3\lambda_4) v'_d v_s/\sqrt{2} + \sqrt{2}\left[-15g_x^2\right] v_d v'_s, \\
M^2_{6,7} &= M^2_{6,8} = -\lambda_5 A_5 v_u/\sqrt{2} + \sqrt{2}\lambda_5^2 v'_d v_s + (\lambda_1\lambda_5 + \lambda_3\lambda_4) v_d v'_s/\sqrt{2} + \sqrt{2}\left[-15g_x^2\right] v'_d v_s, \\
M^2_{6,9} &= -\lambda_3 A_3 v'_u + 2\lambda_3^2 v'_d v'_s + (\lambda_1\lambda_5 + \lambda_3\lambda_4) v_d v_s + 2\left[-15g_x^2\right] v'_d v'_s.
\end{aligned}
\tag{113}
$$

Neutral CP odd Higgs boson

$$
\begin{aligned}
H_a^U &\supset \frac{i\rho_{U,a}}{\sqrt{2}}, \quad H_a^D \supset \frac{i\rho_{D,a}}{\sqrt{2}}, \quad S_a \supset \frac{i\rho_{S,a}}{\sqrt{2}} \quad (a = 1, 2, 3), \\
-\mathcal{L} &\supset \frac{1}{2}\rho_i M^2_{ij} \rho_j, \quad \rho_i = \begin{pmatrix} \rho_{U,a} \\ \rho_{D,a} \\ \rho_{S,a} \end{pmatrix}, \quad (i, j = 1, 2, \cdots, 9),
\end{aligned}
$$

$$
\begin{aligned}
M^2_{1,1} &= M^2_{2,2} = \sqrt{2} m^2_{BU}(v'_u/v_u) + \lambda_1 A_1 v'_s (v_d/v_u) + \lambda_5 A_5 v_s (v'_d/v_u) - \lambda_1^2 v_d^2/2 - \lambda_5^2 v_s^2/2 \\
&- \left[(\lambda_1\lambda_4 + \lambda_3\lambda_5) v_s v'_s + (\lambda_4\lambda_5 + \lambda_1\lambda_3) v'_d v_d\right](v'_u/v_u), \\
M^2_{1,2} &= \lambda_5^2 v_s^2/2 + \lambda_1^2 v_d^2/2, \\
M^2_{1,3} &= M^2_{2,3} = -m^2_{BU} + \left[(\lambda_1\lambda_4 + \lambda_3\lambda_5) v'_s v_s + (\lambda_4\lambda_5 + \lambda_1\lambda_3) v_d v'_d\right]/\sqrt{2}, \\
M^2_{3,3} &= \sqrt{2} m^2_{BU}(v_u/v'_u) + \lambda_3 A_3 v'_s (v'_d/v'_u) + \lambda_4 A_4 v_s (v_d/v'_u) \\
&- \left[(\lambda_1\lambda_4 + \lambda_3\lambda_5) v_s v'_s + (\lambda_4\lambda_5 + \lambda_1\lambda_3) v'_d v_d\right](v_u/v'_u), \\
M^2_{4,4} &= M^2_{5,5} = \sqrt{2} m^2_{BD}(v'_d/v_d) + \lambda_1 A_1 v'_s (v_u/v_d) + \lambda_4 A_4 v_s (v'_u/v_d) - \lambda_1^2 v_u^2/2 - \lambda_4^2 v_s^2/2 \\
&- \left[(\lambda_1\lambda_5 + \lambda_3\lambda_4) v_s v'_s + (\lambda_4\lambda_5 + \lambda_1\lambda_3) v'_u v_u\right](v'_d/v_d), \\
M^2_{4,5} &= \lambda_4^2 v_s^2/2 + \lambda_1^2 v_u^2/2, \\
M^2_{4,6} &= M^2_{5,6} = -m^2_{BD} + \left[(\lambda_1\lambda_5 + \lambda_3\lambda_4) v'_s v_s + (\lambda_4\lambda_5 + \lambda_1\lambda_3) v_u v'_u\right]/\sqrt{2}, \\
M^2_{6,6} &= \sqrt{2} m^2_{BD}(v_d/v'_d) + \lambda_3 A_3 v'_s (v'_u/v'_d) + \lambda_5 A_5 v_s (v_u/v'_d) \\
&- \left[(\lambda_1\lambda_5 + \lambda_3\lambda_4) v_s v'_s + (\lambda_4\lambda_5 + \lambda_1\lambda_3) v'_u v_u\right](v_d/v'_d), \\
M^2_{7,7} &= M^2_{8,8} = \sqrt{2} m^2_{BS}(v'_s/v_s) + \lambda_4 A_4 (v'_u v_d/v_s) + \lambda_5 A_5 (v_u v'_d/v_s) - \lambda_4^2 v_d^2/2 - \lambda_5^2 v_u^2/2 \\
&- \left[(\lambda_1\lambda_5 + \lambda_3\lambda_4) v_d v'_d + (\lambda_1\lambda_4 + \lambda_3\lambda_5) v'_u v_u\right](v'_s/v_s), \\
M^2_{7,8} &= \lambda_4^2 v_d^2/2 + \lambda_5^2 v_u^2/2,
\end{aligned}
$$

$$\begin{aligned}
M^2_{7,9} &= M^2_{8,9} = -m^2_{BS} + [(\lambda_1\lambda_5+\lambda_3\lambda_4)v'_d v_d + (\lambda_1\lambda_4+\lambda_3\lambda_5)v_u v'_u]/\sqrt{2}, \\
M^2_{9,9} &= \sqrt{2}m^2_{BS}(v_s/v'_s) + \lambda_1 A_1(v_u v_d/v'_s) + \lambda_3 A_3(v'_u v'_d/v'_s) \\
&- [(\lambda_1\lambda_5+\lambda_3\lambda_4)v_d v'_d + (\lambda_1\lambda_4+\lambda_3\lambda_5)v'_u v_u]\,(v_s/v'_s), \\
M^2_{1,4} &= M^2_{2,5} = \lambda_1 A_1 v'_s + (\lambda_4\lambda_5-\lambda_1\lambda_3)v'_u v'_d - \lambda_1^2 v_u v_d/2, \\
M^2_{1,5} &= M^2_{2,4} = \lambda_1^2 v_u v_d/2, \\
M^2_{1,6} &= M^2_{2,6} = \lambda_5 A_5 v_s/\sqrt{2} + (\lambda_1\lambda_3-\lambda_4\lambda_5)v'_u v_d/\sqrt{2}, \\
M^2_{3,4} &= M^2_{3,5} = \lambda_4 A_4 v_s/\sqrt{2} + (\lambda_1\lambda_3-\lambda_4\lambda_5)v_u v'_d/\sqrt{2}, \\
M^2_{3,6} &= \lambda_3 A_3 v'_s + (\lambda_4\lambda_5-\lambda_1\lambda_3)v_u v_d, \\
M^2_{1,7} &= M^2_{2,8} = \lambda_5 A_5 v'_d + (\lambda_1\lambda_4-\lambda_3\lambda_5)v'_u v'_s - \lambda_5^2 v_u v_s/2, \\
M^2_{1,8} &= M^2_{2,7} = \lambda_5^2 v_u v_s/2, \\
M^2_{1,9} &= M^2_{2,9} = \lambda_1 A_1 v_d/\sqrt{2} + (\lambda_3\lambda_5-\lambda_1\lambda_4)v'_u v_s/\sqrt{2}, \\
M^2_{3,7} &= M^2_{3,8} = \lambda_4 A_4 v_d/\sqrt{2} + (\lambda_3\lambda_5-\lambda_1\lambda_4)v_u v'_s/\sqrt{2}, \\
M^2_{3,9} &= \lambda_3 A_3 v'_d + (\lambda_1\lambda_4-\lambda_3\lambda_5)v_u v_s, \\
M^2_{4,7} &= M^2_{5,8} = \lambda_4 A_4 v'_u + (\lambda_1\lambda_5-\lambda_3\lambda_4)v'_d v'_s - \lambda_4^2 v_d v_s/2, \\
M^2_{4,8} &= M^2_{5,7} = \lambda_4^2 v_d v_s/2, \\
M^2_{4,9} &= M^2_{5,9} = \lambda_1 A_1 v_u/\sqrt{2} + (\lambda_3\lambda_4-\lambda_1\lambda_5)v'_d v_s/\sqrt{2}, \\
M^2_{6,7} &= M^2_{6,8} = \lambda_5 A_5 v_u/\sqrt{2} + (\lambda_3\lambda_4-\lambda_1\lambda_5)v_d v'_s/\sqrt{2}, \\
M^2_{6,9} &= \lambda_3 A_3 v'_u + (\lambda_1\lambda_5-\lambda_3\lambda_4)v_d v_s.
\end{aligned} \tag{114}$$

Charged Higgs boson

$$\begin{aligned}
&H^U_a \supset H^+_a, \quad H^D_a \supset H^-_{a+3} \quad (a=1,2,3), \\
&-\mathcal{L} \supset H^+_i M^2_{ij} H^-_j, \quad H^+_i = \begin{pmatrix} H^+_a \\ (H^-_{a+3})^\dagger \end{pmatrix} \quad (i,j=1,2,\cdots,6),
\end{aligned}$$

$$\begin{aligned}
M^2_{1,1} &= \sqrt{2}m^2_{BU}(v'_u/v_u) + \lambda_1 A_1 v'_s(v_d/v_u) + \lambda_5 A_5 v_s(v'_d/v_u) - \lambda_1^2 v_d^2 - \lambda_5^2[v_s^2/2 + (v'_d)^2] \\
&- [(\lambda_1\lambda_4+\lambda_3\lambda_5)v_s v'_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v'_d v_d]\,(v'_u/v_u) \\
&- \frac{1}{2}g_2^2[v_u^2 + (v'_u)^2 - v_d^2 - (v'_d)^2] + \frac{1}{4}g_2^2 v_u^2, \\
M^2_{1,2} &= \lambda_5^2 v_s^2/2 + \frac{1}{4}g_2^2 v_u^2, \\
M^2_{1,3} &= M^2_{2,3} = -m^2_{BU} + (\lambda_1\lambda_4+\lambda_3\lambda_5)v_s v'_s/\sqrt{2} + \frac{1}{2\sqrt{2}}g_2^2 v'_u v_u, \\
M^2_{3,3} &= \sqrt{2}m^2_{BU}(v_u/v'_u) + \lambda_3 A_3 v'_s(v'_d/v'_u) + \lambda_4 A_4 v_s(v_d/v'_u) - \lambda_4^2 v_d^2 - \lambda_3^2(v'_d)^2 \\
&- [(\lambda_1\lambda_4+\lambda_3\lambda_5)v_s v'_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v'_d v_d]\,(v_u/v'_u) \\
&- \frac{1}{2}g_2^2[v_u^2 + (v'_u)^2 - v_d^2 - (v'_d)^2] + \frac{1}{2}g_2^2(v'_u)^2, \\
M^2_{4,4} &= M^2_{5,5} = \sqrt{2}m^2_{BD}(v'_d/v_d) + \lambda_1 A_1 v'_s(v_u/v_d) + \lambda_4 A_4 v_s(v'_u/v_d) \\
&- \lambda_1^2 v_u^2 - \lambda_4^2[v_s^2/2 + (v'_u)^2] - [(\lambda_1\lambda_5+\lambda_3\lambda_4)v_s v'_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v'_u v_u]\,(v'_d/v_d) \\
&+ \frac{1}{2}g_2^2[v_u^2 + (v'_u)^2 - v_d^2 - (v'_d)^2] + \frac{1}{4}g_2^2 v_d^2, \\
M^2_{4,5} &= \lambda_4^2 v_s^2/2 + \frac{1}{4}g_2^2 v_d^2, \\
M^2_{4,6} &= M^2_{5,6} = -m^2_{BD} + (\lambda_1\lambda_5+\lambda_3\lambda_4)v_s v'_s/\sqrt{2} + \frac{1}{2\sqrt{2}}g_2^2 v'_d v_d, \\
M^2_{6,6} &= \sqrt{2}m^2_{BD}(v_d/v'_d) + \lambda_3 A_3 v'_s(v'_u/v'_d) + \lambda_5 A_5 v_s(v_u/v'_d) - \lambda_5^2 v_u^2 - \lambda_3^2(v'_u)^2 \\
&- [(\lambda_1\lambda_5+\lambda_3\lambda_4)v_s v'_s + (\lambda_4\lambda_5+\lambda_1\lambda_3)v'_u v_u]\,(v_d/v'_d)
\end{aligned}$$

$$
\begin{aligned}
&+ \quad \frac{1}{2}g_2^2[v_u^2+(v_u')^2-v_d^2-(v_d')^2]+\frac{1}{2}g_2^2(v_d')^2,\\
M_{1,4}^2 &= M_{2,5}^2=\lambda_1A_1v_s'-\lambda_1[\lambda_1v_uv_d+\lambda_3v_u'v_d']+\frac{1}{4}g_2^2v_uv_d,\\
M_{1,5}^2 &= M_{2,4}^2=\frac{1}{4}g_2^2v_uv_d,\\
M_{1,6}^2 &= M_{2,6}^2=\lambda_5A_5v_s/\sqrt{2}-\lambda_5[\lambda_4v_u'v_d+\lambda_5v_uv_d']/\sqrt{2}+\frac{1}{2\sqrt{2}}g_2^2v_uv_d',\\
M_{3,4}^2 &= M_{3,5}^2=\lambda_4A_4v_s/\sqrt{2}-\lambda_4[\lambda_4v_u'v_d+\lambda_5v_uv_d']/\sqrt{2}+\frac{1}{2\sqrt{2}}g_2^2v_u'v_d,\\
M_{3,6}^2 &= \lambda_3A_3v_s'-\lambda_3[\lambda_1v_uv_d+\lambda_3v_u'v_d']+\frac{1}{2}g_2^2v_u'v_d'.
\end{aligned}
\tag{115}
$$

1-loop corrections to Higgs mass

In order to satisfy the experimental bound for the lightest neutral CP even Higgs boson mass, the contributions from 1-loop corrections are important [24][25]. We add 1-loop contributions

$$
\Delta V = \frac{1}{64\pi^2}Str\left[M^4\left(\ln\frac{M^2}{\Lambda^2}-\frac{3}{2}\right)\right]
\tag{116}
$$

to Higgs potentail.The dominant contributions are given by trilinear terms $Y_tH_3^UQ_3U_3^c$ and $kS_3(g_1g_1^c+g_2g_2^c+g_3g_3^c)$. From the mass terms of squark and scalar g-quark

$$
\begin{aligned}
-\mathcal{L} &\supset \left[m_{Q_3}^2+(Y_t(H_3^U)^0)^2\right]|U_3|^2+\left[m_{U_3^c}^2+(Y_t(H_3^U)^0)^2\right]|U_3^c|^2\\
&+ \left[m_g^2+(kv_s')^2\right](|g_1|^2+|g_2|^2+|g_3|^2)+\left[m_{g^c}^2+(kv_s')^2\right](|g_1^c|^2+|g_2^c|^2+|g_3^c|^2)\\
&+ \left[Y_tA_t(H_3^U)^0+Y_t\lambda_3v_s'((H_3^D)^0)^*\right]U_3U_3^c\\
&+ \left[kA_kv_s'+k\lambda_3((H_3^U)^0(H_3^D)^0)^*\right](g_1g_1^c+g_2g_2^c+g_3g_3^c)+h.c.\\
&= (U_3^*,U_3^c)\begin{pmatrix} m_{Q_3}^2+(Y_tv_u')^2 & Y_tA_tv_u'+Y_t\lambda_3v_s'v_d' \\ Y_tA_tv_u'+Y_t\lambda_3v_s'v_d' & m_{U_3^c}^2+(Y_tv_u')^2\end{pmatrix}\begin{pmatrix} U_3 \\ (U_3^c)^*\end{pmatrix}\\
&+ \sum_i(g_i^*,g_i^c)\begin{pmatrix} m_g^2+(kv_s')^2 & kA_kv_s'+k\lambda_3v_u'v_d' \\ kA_kv_s'+k\lambda_3v_u'v_d' & m_{g^c}^2+(kv_s')^2\end{pmatrix}\begin{pmatrix} g_i \\ (g_i^c)^*\end{pmatrix},
\end{aligned}
\tag{117}
$$

mass eigenvalues are given by

$$
\begin{aligned}
M_{T,\pm}^2 &= \frac{1}{2}\left[m_{Q_3}^2+m_{U_3^c}^2+2(Y_tv_u')^2\pm\sqrt{(m_{Q_3}^2-m_{U_3^c}^2)^2+4Y_t^2(A_tv_u'+\lambda_3v_s'v_d')^2}\right],\\
M_{g,\pm}^2 &= \frac{1}{2}\left[m_g^2+m_{g^c}^2+2(kv_s')^2\pm\sqrt{(m_g^2-m_{g^c}^2)^2+4k^2(A_kv_s'+\lambda_3v_u'v_d')^2}\right].
\end{aligned}
\tag{118}
$$

For simplicity, we assume

$$
m_{Q_3}^2=m_{U_3^c}^2=m_g^2=m_{g^c}^2=m_Q^2=16\text{TeV}^2, \quad A_t=A_k=0.0\text{TeV},
\tag{119}
$$

then Eq.(118) is rewritten by

$$
\begin{aligned}
M_{T,\pm}^2 &= m_Q^2+(Y_tv_u')^2\pm Y_t\lambda_3v_s'v_d',\\
M_{g,\pm}^2 &= m_Q^2+(kv_s')^2\pm k\lambda_3v_u'v_d'.
\end{aligned}
\tag{120}
$$

As potential minimum condition is modified as

$$
\frac{\partial(V+\Delta V)}{\partial X}=0, \quad X=(v_u',v_d'),
\tag{121}
$$

and we must add the terms

$$
\begin{aligned}
\Delta M_{3,3}^2 &= \frac{1}{2}\frac{\partial^2\Delta V}{\partial(v_u')^2}-\frac{1}{2(v_u')}\frac{\partial\Delta V}{\partial(v_u')},\\
\Delta M_{6,6}^2 &= \frac{1}{2}\frac{\partial^2\Delta V}{\partial(v_d')^2}-\frac{1}{2(v_d')}\frac{\partial\Delta V}{\partial(v_d')},\\
\Delta M_{3,6}^2 &= \frac{1}{2}\frac{\partial^2\Delta V}{\partial(v_u')\partial(v_d')},
\end{aligned}
\tag{122}
$$

to the neutral CP-even Higgs boson mass. We fix renormalization point as

$$\Lambda = 4\mathrm{TeV}. \tag{123}$$

Chargino

$$\begin{aligned}
\mathcal{L} &\supset \chi^+_i M_{ij} \chi^-_j + h.c., \\
& h^U_a = \begin{pmatrix} h^+_u \\ h^0_u \end{pmatrix}_a, \quad h^D_a = \begin{pmatrix} h^0_d \\ h^-_d \end{pmatrix}_a \quad (a = 1, 2, 3), \\
& \chi^+_i = \begin{pmatrix} (h^+_u)_a \\ -iw^+ \end{pmatrix}, \quad \chi^-_i = \begin{pmatrix} (h^-_d)_a \\ -iw^- \end{pmatrix}, \quad w^\pm = \frac{\lambda^1_2 \mp i\lambda^2_2}{\sqrt{2}} \quad (i, j = 1, 2, 3, 4), \\
M &= \begin{pmatrix} \lambda_1 v'_s & 0 & \lambda_5 v_s/\sqrt{2} & g_2 v_u/\sqrt{2} \\ 0 & \lambda_1 v'_s & \lambda_5 v_s/\sqrt{2} & g_2 v_u/\sqrt{2} \\ \lambda_4 v_s/\sqrt{2} & \lambda_4 v_s/\sqrt{2} & \lambda_3 v'_s & g_2 v'_u \\ g_2 v_d/\sqrt{2} & g_2 v_d/\sqrt{2} & g_2 v'_d & M_2 \end{pmatrix}.
\end{aligned} \tag{124}$$

Neutralino

$$\begin{aligned}
& \mathcal{L} \supset -\frac{1}{2} \eta_i M_{ij} \eta_j + h.c., \\
& \eta = \begin{pmatrix} (h^0_u)_a \\ (h^0_d)_a \\ s_a \\ i\lambda \end{pmatrix}, \quad \lambda = \begin{pmatrix} \lambda_Y \\ \lambda^3_2 \\ \lambda_X \end{pmatrix} \quad (a = 1, 2, 3; \quad i, j = 1, 2, \cdots, 12), \\
& M = \begin{pmatrix} 0 & M_{ud} & M_{us} & M_{u\lambda} \\ M_{du} & 0 & M_{ds} & M_{d\lambda} \\ M_{su} & M_{sd} & 0 & M_{s\lambda} \\ M_{\lambda u} & M_{\lambda d} & M_{\lambda s} & M_{\lambda\lambda} \end{pmatrix},
\end{aligned}$$

$$\begin{aligned}
M_{ud} &= \begin{pmatrix} \lambda_1 v'_s & 0 & \lambda_5 v_s/\sqrt{2} \\ 0 & \lambda_1 v'_s & \lambda_5 v_s/\sqrt{2} \\ \lambda_4 v_s/\sqrt{2} & \lambda_4 v_s/\sqrt{2} & \lambda_3 v'_s \end{pmatrix} = M^T_{du}, \\
M_{us} &= \begin{pmatrix} \lambda_5 v'_d & 0 & \lambda_1 v_d/\sqrt{2} \\ 0 & \lambda_5 v'_d & \lambda_1 v_d/\sqrt{2} \\ \lambda_4 v_d/\sqrt{2} & \lambda_4 v_d/\sqrt{2} & \lambda_3 v'_d \end{pmatrix} = M^T_{su}, \\
M_{ds} &= \begin{pmatrix} \lambda_4 v'_u & 0 & \lambda_1 v_u/\sqrt{2} \\ 0 & \lambda_4 v'_u & \lambda_1 v_u/\sqrt{2} \\ \lambda_5 v_u/\sqrt{2} & \lambda_5 v_u/\sqrt{2} & \lambda_3 v'_u \end{pmatrix} = M^T_{sd}, \\
M_{u\lambda} &= \begin{pmatrix} g_Y v_u/2 & -g_2 v_u/2 & -2g_x v_u \\ g_Y v_u/2 & -g_2 v_u/2 & -2g_x v_u \\ g_Y v'_u/\sqrt{2} & -g_2 v'_u/\sqrt{2} & -2\sqrt{2} g_x v'_u \end{pmatrix} = M^T_{\lambda u}, \\
M_{d\lambda} &= \begin{pmatrix} -g_Y v_d/2 & g_2 v_d/2 & -3g_x v_d \\ -g_Y v_d/2 & g_2 v_d/2 & -3g_x v_d \\ -g_Y v'_d/\sqrt{2} & g_2 v'_d/\sqrt{2} & -3\sqrt{2} g_x v'_d \end{pmatrix} = M^T_{\lambda d}, \\
M_{s\lambda} &= \begin{pmatrix} 0 & 0 & 5g_x v_s \\ 0 & 0 & 5g_x v_s \\ 0 & 0 & 5\sqrt{2} g_x v'_s \end{pmatrix} = M^T_{\lambda s}, \\
M_{\lambda\lambda} &= \begin{pmatrix} -M_Y & 0 & 0 \\ 0 & -M_2 & 0 \\ 0 & 0 & -M_X \end{pmatrix}.
\end{aligned} \tag{125}$$

B Mixing matrices and mass eigenvalues

Mass eigenvalues (m_i : GeV) and diagnalization matrix $U = (u_1, u_2, \cdots)$ are given as follows.

Neutral CP-even Higgs

For Higgs bosons, the diagonalization matrices are defined as

$$(M^2)' = U^T M^2 U = \mathrm{diag}(m_1^2, m_2^2, \cdots). \tag{126}$$

i	1	2	3	4	5	6	7	8	9
u_i	−0.698	−0.113	−0.0006	−0.613	0.329	−0.115	0.0006	0.003	−0.059
	0.698	0.113	0.0006	−0.613	0.329	−0.115	0.0006	0.003	−0.059
	0	0	0	−0.384	−0.808	−0.036	0.029	−0.009	−0.445
	0.0015	−0.005	−0.707	−0.0012	−0.008	0.006	−0.706	−0.0008	−0.030
	−0.0015	0.005	0.707	−0.0012	−0.008	0.006	−0.706	−0.0008	−0.030
	0	0	0	−0.273	−0.360	−0.031	−0.033	0.054	0.889
	0.113	−0.698	−0.0056	0.118	−0.025	−0.693	0.006	0.071	−0.002
	−0.113	0.698	0.0056	0.118	−0.025	−0.693	0.006	0.071	−0.002
	0	0	0	−0.002	0.014	0.101	0.00009	0.993	−0.052
m_i	91.50	537	2005	121.96	152.96	539	2007	1018	1425

(127)

Neutral CP-odd Higgs

i	1	2	3	4	5	6	7	8	9
u_i	−0.703	0.073	−0.040	0.004	0.0005	0.700	0.074	−0.002	−0.059
	0.703	−0.073	−0.040	0.004	−0.0005	0.700	0.074	−0.002	−0.059
	0	0	−0.889	0.086	0	−0.088	−0.009	−0.029	−0.044
	0.0014	0.008	0.008	−0.00008	0.707	0.0003	0.008	−0.706	0.030
	−0.0014	0.008	0.008	−0.00008	−0.707	0.0003	0.008	−0.706	0.030
	0	0	0.448	−0.004	0	−0.049	−0.005	−0.033	−0.892
	−0.073	−0.703	−0.006	−0.070	0.008	0.074	−0.700	−0.008	−0.002
	0.073	0.703	−0.006	−0.070	−0.008	0.074	−0.700	−0.008	−0.002
	0	0	−0.079	−0.991	0	−0.012	0.099	−0.001	−0.034
m_i	112.12	532	0	0	2005	145.53	535	2008	1423

(128)

Charged Higgs

i	1	2	3	4	5	6
u_i	0.707	−0.000055	−0.703	0.041	−0.001	0.059
	−0.707	0.000055	−0.703	0.041	−0.001	0.059
	0	0	0.089	0.893	−0.029	0.441
	0.000055	0.707	−0.0018	−0.008	−0.706	−0.030
	−0.000055	−0.707	−0.0018	−0.008	−0.706	−0.030
	0	0	0.049	−0.447	−0.032	0.893
m_i	90.70	2005	130.02	0	2007	1422

(129)

Neutralino

For neutralinos, the diagonalization matrix is defined as

$$U^T M U = \mathrm{diag}(m_1, m_2, \cdots m_{12}).$$

i	1	2	3	4	5	6
u_i	0.234	-0.496	0.446	-0.248	-0.475	0.067
	-0.234	0.496	-0.446	-0.248	-0.475	0.067
	0	0	0	0.064	0.148	0.031
	0.209	0.504	0.450	-0.217	0.497	-0.055
	-0.209	-0.504	-0.450	-0.217	0.497	-0.055
	0	0	0	0.028	-0.105	-0.066
	-0.634	-0.017	0.313	0.621	-0.023	-0.002
	0.634	0.017	-0.313	0.621	-0.023	-0.002
	0	0	0	-0.086	0.005	-0.00007
	0	0	0	0.009	0.084	0.965
	0	0	0	-0.014	-0.119	-0.221
	0	0	0	0.0004	0.001	0.003
m_i	-41.92	-130	172	-44.55	-111	-177

i	7	8	9	10	11	12
u_i	0.062	-0.430	-0.099	-0.099	-0.020	0.002
	0.062	-0.430	-0.099	-0.099	-0.020	0.002
	0.039	0.132	-0.689	-0.682	-0.121	-0.003
	-0.049	-0.439	0.062	-0.063	-0.013	-0.003
	-0.049	-0.439	0.062	-0.063	-0.013	-0.003
	-0.077	0.085	0.692	-0.686	-0.138	-0.050
	-0.002	-0.330	0.001	-0.003	-0.053	0.047
	-0.002	-0.330	0.001	-0.003	-0.053	0.047
	-0.0001	0.051	0.014	0.114	-0.728	0.669
	-0.241	0.007	0.063	-0.014	-0.002	-0.001
	-0.960	-0.012	-0.118	0.024	0.003	0.002
	0.003	-0.0009	-0.032	0.153	-0.655	-0.739
m_i	-194	157	832	820	945	-1144

(130)

Chargino
For the charginos, the diagonalization matrices are defined as

$$\begin{aligned}
&\chi^- = U_-(\chi^-)', \quad \chi^+ = U_+(\chi^+)', \\
&U_+^T M U_- = diag(m_1, m_2, m_3, m_4) = (130, 111, 193, 830), \\
&U_+ = \begin{pmatrix} 0.707 & 0.685 & 0.104 & 0.140 \\ -0.707 & 0.685 & 0.104 & 0.140 \\ 0 & -0.209 & 0.066 & 0.976 \\ 0 & 0.131 & -0.987 & 0.095 \end{pmatrix}, \quad U_- = \begin{pmatrix} -0.707 & -0.696 & 0.085 & 0.088 \\ 0.707 & -0.696 & 0.085 & 0.088 \\ 0 & 0.140 & 0.128 & 0.982 \\ 0 & -0.102 & -0.984 & 0.143 \end{pmatrix}.
\end{aligned} \tag{131}$$

These mass eigenvalues are consistent with the experimental mass bounds [18]

Charged Higgs : $m \geq 79.3$GeV,
Neutral CP-even Higgs : $m \geq 114.4$GeV,
Neutral CP-odd Higgs : $m \geq 93.4$GeV,
Chargino : $m \geq 94$GeV,
Neutralino : $m \geq 46$GeV.

Note that $\rho_3', \rho_4', (H_4^\pm)'$ are Nambu-Goldstone boson which are eaten by gauge bosons.

C The lifetimes of exotic quarks

As the R-parities of exotic quarks are odd, at least there must be one sfermion which is lighter than exotic quarks, to make them unstable. Now we assume the right handed slepton E_1^c is lighter than $2TeV$ and the other sfermions are heavier than $4TeV$. For simplicity, we assume there is no mixing between E_1^c and L_i.

Through the non-renormalizable interaction

$$\mathcal{L} \supset -\sum_i \frac{c_i V^2}{\sqrt{3}M_P^2}(\psi_{g,1}+\psi_{g,2}+\psi_{g,3})E_1^c u_i^c + h.c., \tag{132}$$

the exotic quarks $\psi_{g,1\sim 3}$ can decay into u_i^c and E_1^c, where c_i are $O(1)$ coefficients. The lifetimes are estimated as follows

$$\Gamma(\psi_{g,j}^\dagger \to E_1^c + u_i^c) = \frac{2TeV}{16\pi}\left(\frac{c_i V^2}{\sqrt{3}M_P^2}\right)^2 = \frac{c_i^2}{1.7[\mathrm{sec}]}, \tag{133}$$

from which we must put $c_i \sim 4$ in order to satisfy the cosmological constraint for exotic particle, $\tau < 0.1$sec. The interaction Eq.(132) comes from

$$W \supset \sum_i \frac{c_i}{M_P^2}\Phi_3(\Phi_1^c g_1 + \Phi_2^c g_2 + \Phi_3^c g_3)E_1^c U_i^c, \tag{134}$$

which may contribute to LF decay through

$$\mathcal{L} \supset -\sum_{i,j} \frac{c_i V}{M_P^2}\alpha_j \phi_{LF}\psi_{g,j}E_1^c u_i^c + h.c., \tag{135}$$

where α_j are given by linear combinations of the flavon mixing parameters. In this paper, we assume α_j are small enough and this interaction does not give sizable contribution to LF decay width.

Acknowledgments

This work is supported by Young Researcher Overseas Visits Program for Vitalizing Brain Circulation Japanese in JSPS (T.T.). H.O. acknowledges partial supports from the Science and Technology Development Fund (STDF) project ID 437 and the ICTP project ID 30.

References

[1] O. Adriani *et al.* [PAMELA Collaboration], Nature **458** (2009) 607.

[2] P. Meade, M. Papucci, A. Strumia and T. Volansky, Nucl. Phys. **B831** (2010) 178; A. Ibarra, D. Tran and C. Weniger, JCAP**1001** (2010) 009; C. R. Chen, S. K. Mandal and F. Takahashi, JCAP**1001** (2010) 023; J. Liu, Q. Yuan, X. Bi, H. Li and X. Zhang, arXiv:0911.1002; M. Cirelli, P. Panci and P. D. Serpico, arXiv:0912.0663; G. Hutsi, A. Hektor and M. Raidal, arXiv:1004.2036; A. Ibarra and D. Tran, JCAP**0807** (2008) 002.

[3] A. Ibarra and D. Tran, JCAP **0902**, 021 (2009) [arXiv:0811.1555 [hep-ph]].

[4] G. Bertone, M. Cirelli, A. Strumia and M. Taoso, JCAP **03** (2009) 009; M. Cirelli and P. Panci, Nucl. Phys. **B821**, (2009) 399; M. Papucci and A. Strumia, JCAP **03** (2010) 014.

[5] H.P. Nilles, Phys. Rep. **110** (1984) 1.

[6] H. Murayama and J. D. Wells, Phys. Rev. **D65** (2002) 055009.

[7] J. Kubo, Phys. Lett. **B578** (2004) 156.

[8] Y. Daikoku and H. Okada, Phys. Rev. D **82**, 033007 (2010).

[9] Y. Kajiyama, J. Kubo and H. Okada, Phys. Rev. D **75**, 033001 (2007) [arXiv:hep-ph/0610072]; Y. Kajiyama, H. Okada and T. Toma, arXiv:1104.0367 [hep-ph].

[10] See for example the reviews of Non-Abelian discrete flavor symmetries by G. Altarelli, F. Feruglio, Rev. Mod. Phys. **82**, 2701-2729 (2010). [arXiv:1002.0211 [hep-ph]]; H. Ishimori, T. Kobayashi, H. Ohki, H. Okada, Y. Shimizu and M. Tanimoto, Prog. Theor. Phys. Suppl. **183**, 1 (2010);

[11] S. Pakvasa and H. Sugawara, Phys. Lett. **B73** (1978) 61; E. Ma, Phys. Lett. **B632** (2006) 352; C. Hagedorn, M. Lindner and R.N. Mohapatra, JHEP**0606** (2006) 042; H. Zhang, Phys. Lett. B **655**, 132 (2007); Y. Koide, JHEP**0708** (2007) 086; F. Bazzocchi and S. Morisi, Phys. Rev. D **80**, 096005 (2009); H. Ishimori, Y. Shimizu and M. Tanimoto, Prog. Theor. Phys. **121**, 769 (2009); F. Bazzocchi, L. Merlo and S. Morisi, Nucl. Phys. B **816** (2009) 204; F. Bazzocchi, L. Merlo and S. Morisi, Phys. Rev. D **80** (2009) 053003 G. Altarelli, F. Feruglio and L. Merlo, JHEP **0905** (2009) 020; W. Grimus, L. Lavoura and P. O. Ludl, J. Phys. G **36**, 115007 (2009); G. J. Ding, Nucl. Phys. B **827**, 82 (2010); B. Dutta, Y. Mimura and R. N. Mohapatra, JHEP **1005**, 034 (2010); D. Meloni, J. Phys. G **37**, 055201 (2010); S. Morisi and E. Peinado, Phys. Rev. D **81**, 085015 (2010); C. Hagedorn, S. F. King and C. Luhn, JHEP **1006**, 048 (2010); R. Z. Yang and H. Zhang, Phys. Lett. B **700**, 316 (2011) [arXiv:1104.0380 [hep-ph]]; Y. Shimizu, M. Tanimoto and A. Watanabe, arXiv:1105.2929 [hep-ph]; H. Ishimori, Y. Shimizu, M. Tanimoto and A. Watanabe, Phys. Rev. D **83**, 033004 (2011) [arXiv:1010.3805 [hep-ph]]; H. Ishimori, K. Saga, Y. Shimizu and M. Tanimoto, Phys. Rev. D **81**, 115009 (2010) [arXiv:1004.5004 [hep-ph]]; K. M. Patel, Phys. Lett. B **695**, 225 (2011) [arXiv:1008.5061 [hep-ph]].

[12] D. Suematsu and Y. Yamagishi, Int. J. Mod. Phys. **A10** (1995) 4521.

[13] Y. Daikoku, H. Okada and T. Toma [arXiv:1010.4963 [hep-ph]].

[14] R. Howl and S.F. King, JHEP**0805** (2008) 008.

[15] K. Abe *et al.* [T2K Collaboration], arXiv:1106.2822 [hep-ex].

[16] N. Haba, Y. Kajiyama, S. Matsumoto, H. Okada and K. Yoshioka, Phys. Lett. B **695**, 476 (2011) [arXiv:1008.4777 [hep-ph]]; Y. Kajiyama and H. Okada, Nucl. Phys. B **848**, 303 (2011) [arXiv:1011.5753 [hep-ph]].

[17] J. Kubo, H. Okada and F. Sakamaki, Phys. Rev. D **70**, 036007 (2004) [arXiv:hep-ph/0402089]; Y. Daikoku and H. Okada, arXiv:1008.0914 [hep-ph].

[18] C. Amsler et al. (Particle Data Group), Phys. Lett. B667, 1 (2008) and 2009 partial update for the 2010 edition. Cut-off date for this update was January 15, 2009.

[19] Z. z. Xing, H. Zhang and S. Zhou, Phys. Rev. **D77** (2008) 113016.

[20] F. Riva, Phys. Lett. B **690**, 443 (2010) [arXiv:1004.1177 [hep-ph]].

[21] S. Matsumoto and K. Yoshioka, Phys. Rev. **D82** (2010) 053009.

[22] M. Kawasaki, K. Kohri and T. Moroi, Phys. Rev. **D71** (2005) 083502.

[23] O. Adriani *et al.*, Phys. Rev. Lett. **102** (2009) 051101.

[24] Y. Okada, M. Yamaguchi and T. Yanagida, Prog. Theor. Phys. **85** (1991) 1.

[25] Y. Daikoku and D. Suematsu, Prog. Theor. Phys. **104** (2000) 104, Y. Daikoku and D. Suematsu, Phys. Rev. **D62** (2000) 095006.

[26] Q. H. Cao, E. Ma, J. Wudka and C. P. Yuan, arXiv:0711.3881 [hep-ph]; M. Fairbairn and J. Zupan, JCAP **0907**, 001 (2009) [arXiv:0810.4147 [hep-ph]]; H. Fukuoka, D. Suematsu and T. Toma, arXiv:1012.4007 [hep-ph]; D. Suematsu and T. Toma, Nucl. Phys. B **847**, 567 (2011) [arXiv:1011.2839 [hep-ph]].

[27] T. Sjostrand, S. Mrenna and P. Z. Skands, JHEP **0605**, 026 (2006) [arXiv:hep-ph/0603175].